

Sums, series, and products in Diophantine approximation

Jordan Bell

jordan.bell@gmail.com

Department of Mathematics, University of Toronto
Toronto, Ontario, Canada

June 18, 2023

Abstract

There is not much that can be said for all x and for all n about the sum

$$\sum_{k=1}^n \frac{1}{|\sin k\pi x|}.$$

However, for this and similar sums, series, and products, we can establish results for almost all x using the tools of continued fractions. We present in detail the appearance of these sums in the singular series for the circle method. One particular interest of the paper is the detailed proof of a striking result of Hardy and Littlewood, whose compact proof, which delicately uses analytic continuation, has not been written freshly anywhere since its original publication. This story includes various parts of late 19th century and early 20th century mathematics.

Contents

1	Introduction	2
2	The Bernoulli polynomials, the Euler-Maclaurin summation formula, and Euler's constant	3
3	Background	7
4	Preliminaries on continued fractions	12
5	Diophantine conditions	17
6	Sums of reciprocals	19
7	Weyl's inequality, Vinogradov's estimate, Farey fractions, and the circle method	30

8	Discrepancy and exponential sums	40
9	Dirichlet series	47
10	Power series	49
11	Product	53
12	Conclusions	68

1 Introduction

In this paper we survey a class of estimates for sums, series, and products that involve Diophantine approximation. We both sort out a timeline of the literature on these questions and give careful proofs of many lesser known results. Rather than being an open-pit mine for the history of Diophantine approximation, this paper follows one vein as deep as it goes.

For $x \in \mathbb{R}$, we write $\|x\| = \min_{n \in \mathbb{Z}} |x - n|$, the distance from x to a nearest integer. In this paper we give a comprehensive presentation of estimates for sums whose j th term involves $\|jx\|$ and determine the abscissa of convergence and radius of convergence respectively of Dirichlet series and power series whose j th term involves $\|jx\|$. We also give a detailed proof of a result of Hardy and Littlewood that

$$\lim_{n \rightarrow \infty} \left(\prod_{k=1}^n |\sin k\pi x| \right)^{1/n} = \frac{1}{2}$$

for almost all x .

We either prove or state in detail and give references for all the material on continued fractions and measure theory that we use in this paper. Many of the results we prove in this paper do not have detailed proofs written in any books, and the proofs we give for results that do have proofs in books are often written significantly more meticulously here than anywhere else; in some cases the proofs in the literature are so sketchy that the proof we give is written from scratch, for example Lemma 9. Our presentation of Hardy and Littlewood's estimate for $\prod_{k=1}^n |\sin \pi kx|$ makes clear exactly what results in Diophantine approximation one needs for the proof.

In the next section we introduce the Bernoulli polynomials, the Euler-Maclaurin summation formula, and Euler's constant, which we shall use in a few places. Because the calculations are similar to what we do in the rest of this paper and because we want to be comfortable using Bernoulli polynomials, we work things out from scratch rather than merely stating results as known. In the section after that we summarize various problems that involve sums of the type we are talking about in this paper.

2 The Bernoulli polynomials, the Euler-Maclaurin summation formula, and Euler's constant

For $k \geq 0$, the **Bernoulli polynomial** $B_k(x)$ is defined by

$$\frac{ze^{xz}}{e^z - 1} = \sum_{k=0}^{\infty} B_k(x) \frac{z^k}{k!}, \quad |z| < 2\pi. \quad (1)$$

The **Bernoulli numbers** are $B_k = B_k(0)$, the constant terms of the Bernoulli polynomials. For any x , using L'Hospital's rule the left-hand side of (1) tends to 1 as $z \rightarrow 0$, and the right-hand side tends to $B_0(x)$, hence $B_0(x) = 1$. Differentiating (1) with respect to x ,

$$\sum_{k=0}^{\infty} B'_k(x) \frac{z^k}{k!} = \frac{z^2 e^{xz}}{e^z - 1} = \sum_{k=0}^{\infty} B_k(x) \frac{z^{k+1}}{k!} = \sum_{k=1}^{\infty} B_{k-1}(x) \frac{z^k}{(k-1)!},$$

so $B'_0(x) = 0$ and for $k \geq 1$ we have $\frac{B'_k(x)}{k!} = \frac{B_{k-1}(x)}{(k-1)!}$, i.e.

$$B'_k(x) = kB_{k-1}(x).$$

Furthermore, integrating (1) with respect to x on $[0, 1]$ gives, since $\int e^{xz} dx = \frac{e^z - 1}{z}$,

$$1 = \sum_{k=0}^{\infty} \left(\int_0^1 B_k(x) dx \right) \frac{z^k}{k!}, \quad |z| < 2\pi,$$

hence $\int_0^1 B_0(x) dx = 1$ and for $k \geq 1$,

$$\int_0^1 B_k(x) dx = 0.$$

The first few Bernoulli polynomials are

$$B_0(x) = 1, \quad B_1(x) = x - \frac{1}{2}, \quad B_2(x) = x^2 - x + \frac{1}{6}, \quad B_3(x) = x^3 - \frac{3}{2}x^2 + \frac{1}{2}x.$$

The Bernoulli polynomials satisfy the following:

$$\begin{aligned} \sum_{k=0}^{\infty} B_k(x+1) \frac{z^k}{k!} &= \frac{ze^{(x+1)z}}{e^z - 1} \\ &= \frac{ze^{xz}(e^z - 1 + 1)}{e^z - 1} \\ &= ze^{xz} + \frac{ze^{xz}}{e^z - 1} \\ &= \sum_{k=0}^{\infty} \frac{x^k z^{k+1}}{k!} + \sum_{k=0}^{\infty} B_k(x) \frac{z^k}{k!} \\ &= \sum_{k=1}^{\infty} \frac{x^{k-1} z^k}{(k-1)!} + \sum_{k=0}^{\infty} B_k(x) \frac{z^k}{k!}, \end{aligned}$$

hence

$$B_k(x+1) = kx^{k-1} + B_k(x), \quad k \geq 1, x \in \mathbb{R}. \quad (2)$$

In particular, for $k \geq 2$, $B_k(1) = B_k(0)$. The identity (2) yields **Faulhaber's formula**, for $k \geq 0$ and positive integers $a < b$,

$$\sum_{a \leq m \leq b} m^k = \frac{B_{k+1}(b+1) - B_{k+1}(a)}{k+1}. \quad (3)$$

The following identity is the **multiplication formula for the Bernoulli polynomials**, found by Raabe [122, pp. 19-24, §13].

Lemma 1. For $k \geq 0$, $q \geq 1$, and $x \in \mathbb{R}$,

$$qB_k(qx) = q^k \sum_{j=0}^{q-1} B_k\left(x + \frac{j}{q}\right).$$

Proof. Using (2) with $x = \frac{qn+j}{q} = n + \frac{j}{q}$,

$$(qn+j)^k = \frac{q^k}{k+1} \left(B_{k+1}\left(n + \frac{j}{q} + 1\right) - B_{k+1}\left(n + \frac{j}{q}\right) \right),$$

thus

$$\begin{aligned} \sum_{m=q}^{Nq-1} m^k &= \sum_{n=1}^{N-1} \sum_{j=0}^{q-1} (nq+j)^k \\ &= \frac{q^k}{k+1} \sum_{j=0}^{q-1} \sum_{n=1}^{N-1} \left(B_{k+1}\left(n + \frac{j}{q} + 1\right) - B_{k+1}\left(n + \frac{j}{q}\right) \right) \\ &= \frac{q^k}{k+1} \sum_{j=0}^{q-1} \left(B_{k+1}\left(N + \frac{j}{q}\right) - B_{k+1}\left(1 + \frac{j}{q}\right) \right). \end{aligned}$$

Then by (3),

$$B_{k+1}(qN) - B_{k+1}(q) = q^k \sum_{j=0}^{q-1} \left(B_{k+1}\left(N + \frac{j}{q}\right) - B_{k+1}\left(1 + \frac{j}{q}\right) \right).$$

Let $F_{k,q}(x) = B_{k+1}(qx) - q^k \sum_{j=0}^{q-1} B_{k+1}(x + j/q)$, which is a polynomial of degree $\leq k+1$. The above means that for $N \geq 1$, $F_{k,q}(N) = F_{k,q}(1)$, which implies that the polynomial $F_{k,q}(x) - F_{k,q}(1)$ is identically 0 and, a fortiori, $F'_{k,q}(x)$ is identically 0:

$$qB'_{k+1}(qx) - q^k \sum_{j=0}^{q-1} B'_{k+1}\left(x + \frac{j}{q}\right) = 0.$$

Using $B'_{k+1}(x) = (k+1)B_k(x)$,

$$qB_k(qx) = q^k \sum_{j=0}^{q-1} B_k\left(x + \frac{j}{q}\right).$$

□

We remark that for prime p and for $k \geq 0$, one uses Lemma 1 to prove that there is a unique p -adic distribution $\mu_{B,k}$ on the p -adic integers \mathbb{Z}_p such that $\mu_{B,k}(a + p^N \mathbb{Z}_p) = p^{N(k-1)} B_k(a/p^N)$ [80, p. 35, Chapter II, §4], called a **Bernoulli distribution**.

For $x \in \mathbb{R}$, let $[x]$ be the greatest integer $\leq x$, and let $R(x) = x - [x]$, called the **fractional part of x** . Write $\mathbb{T} = \mathbb{R}/\mathbb{Z}$ and define the **periodic Bernoulli functions** $P_k : \mathbb{T} \rightarrow \mathbb{R}$ by

$$P_k = B_k \circ R.$$

For $k \geq 2$, because $B_k(1) = B_k(0)$, the function P_k is continuous. For $f : \mathbb{T} \rightarrow \mathbb{C}$ define its **Fourier series** $\hat{f} : \mathbb{Z} \rightarrow \mathbb{C}$ by

$$\hat{f}(n) = \int_{\mathbb{T}} f(t) e^{-2\pi i n t} dt, \quad n \in \mathbb{Z}.$$

For $k \geq 1$, one calculates $\hat{P}_k(0) = 0$ and using integration by parts, $\hat{P}_k(n) = -\frac{1}{(2\pi i)^k}$ for $n \neq 0$. Thus for $k \geq 1$, the Fourier series of P_k is

$$P_k(t) \sim \sum_{n \in \mathbb{Z}} \hat{P}_k(n) e^{2\pi i n t} = -\frac{1}{(2\pi i)^k} \sum_{n \neq 0} n^{-k} e^{2\pi i n t}.$$

For $k \geq 2$, $\sum_{n \in \mathbb{Z}} |\hat{P}_k(n)| < \infty$, from which it follows that $\sum_{|n| \leq N} \hat{P}_k(n) e^{2\pi i n t}$ converges to $P_k(t)$ uniformly for $t \in \mathbb{T}$. Furthermore, for $t \notin \mathbb{Z}$ [103, p. 499, Theorem B.2],

$$P_1(t) = -\frac{1}{\pi} \sum_{n=1}^{\infty} \frac{1}{n} \sin 2\pi n t.$$

Thus for example,

$$B_1\left(\frac{1}{2\pi}\right) = P_1\left(\frac{1}{2\pi}\right) = -\frac{1}{\pi} \sum_{k=1}^{\infty} \frac{\sin k}{k}.$$

The **Euler-Maclaurin summation formula** is the following [103, p. 500, Theorem B.5]. If $a < b$ are real numbers, K is a positive integer, and f is a C^K function on an open set that contains $[a, b]$, then

$$\begin{aligned} \sum_{a < m \leq b} f(m) &= \int_a^b f(x) dx + \sum_{k=1}^K \frac{(-1)^k}{k!} (P_k(b) f^{(k-1)}(b) - P_k(a) f^{(k-1)}(a)) \\ &\quad - \frac{(-1)^K}{K!} \int_a^b P_K(x) f^{(K)}(x) dx. \end{aligned}$$

Applying the Euler-Maclaurin summation formula with $a = 1, b = n, K = 2, f(x) = \log x$ yields [103, p. 503, Eq. B.25]

$$\sum_{1 \leq m \leq n} \log m = n \log n - n + \frac{1}{2} \log n + \frac{1}{2} \log 2\pi + O(n^{-1}).$$

Using $e^{1+O(n^{-1})} = 1 + O(n^{-1})$, taking the exponential of the above equation gives **Stirling's approximation**,

$$n! = n^n e^{-n} \sqrt{2\pi n} (1 + O(n^{-1})).$$

Write $a_n = -\log n + \sum_{1 \leq m \leq n} \frac{1}{m}$. Because $x \mapsto \log(1-x)$ is concave,

$$a_n - a_{n-1} = \frac{1}{n} + \log\left(1 - \frac{1}{n}\right) \leq 1 + 1 - \frac{1}{n} = 0,$$

which means that the sequence a_n is nonincreasing. For $f(x) = \frac{1}{x}$, because f is positive and nonincreasing,

$$\sum_{1 \leq m \leq n} f(m) \geq \int_1^{n+1} f(x) dx = \log(n+1) > \log n,$$

hence $a_n > 0$. Because the sequence a_n is positive and nonincreasing, there exists some nonnegative limit, γ , called **Euler's constant**. Using the Euler-Maclaurin summation formula with $a = 1, b = n, K = 1, f(x) = \frac{1}{x}$, as $P_1(x) = [x] - \frac{1}{2}$,

$$\sum_{1 < m \leq n} \frac{1}{m} = \log n + \frac{1}{2n} - \frac{1}{2} + \frac{1}{2} \int_1^n \frac{1}{x^2} dx - \int_1^n R(x) \frac{1}{x^2} dx,$$

which is

$$\sum_{1 < m \leq n} \frac{1}{m} = \log n - \int_1^\infty \frac{R(x)}{x^2} dx + \int_n^\infty \frac{R(x)}{x^2} dx.$$

As $0 \leq R(x)x^{-2} \leq x^{-2}$, the function $x \mapsto R(x)x^{-2}$ is integrable on $[1, \infty)$; let $C = 1 - \int_1^\infty R(x)x^{-2}$. Since $0 \leq \int_n^\infty R(x)x^{-2} dx \leq \int_n^\infty x^{-2} dx = n^{-1}$,

$$\sum_{1 \leq m \leq n} \frac{1}{m} = \log n + C + O(n^{-1})$$

f. But $-\log n + \sum_{1 \leq m \leq n} \frac{1}{m} \rightarrow \gamma$ as $n \rightarrow \infty$, from which it follows that $C = \gamma$ and thus

$$\sum_{1 \leq m \leq n} \frac{1}{m} = \log n + \gamma + O(n^{-1}).$$

3 Background

For $x \in \mathbb{R}$, let $[x]$ be the greatest integer $\leq x$ and let $R(x) = x - [x]$. It will be handy to review some properties of $x \mapsto [x]$. For $n \in \mathbb{Z}$ it is immediate that $[x + n] = [x]$. For $x, y \in \mathbb{R}$ we have $0 \leq x - [x] + y - [y] < 2$, which means $0 = [0] \leq [x - [y] + y - [y]] < [2] = 2$, and using $[x + n] = [x]$ this is $0 \leq [x + y] - [x] - [y] < 2$, therefore

$$[x] + [y] \leq [x + y] \leq [x] + [y] + 1.$$

For $m, n \in \mathbb{Z}$, $n > 1$, and $x \in \mathbb{R}$,

$$\left[\frac{x + m}{n} \right] = \left[\frac{[x] + m}{n} \right].$$

For n a positive integer and for real x ,

$$[x] = \left[x + \frac{0}{n} \right] \leq \left[x + \frac{1}{n} \right] \leq \cdots \leq \left[x + \frac{n-1}{n} \right] \leq \left[x + \frac{n}{n} \right] = [x] + 1.$$

There is a unique ν , $1 \leq \nu \leq n$, such that $[x + \frac{\nu-1}{n}] = [x]$ and $[x + \frac{\nu}{n}] = [x] + 1$, and therefore $[R(x) + \frac{\nu-1}{n}] = 0$ and $[R(x) + \frac{\nu}{n}] = 1$, consequently $R(x) + \frac{\nu-1}{n} < 1$ and $R(x) + \frac{\nu}{n} \geq 1$, which means $1 - \frac{\nu}{n} \leq R(x) < 1 - \frac{\nu-1}{n}$, from which finally we get $n \leq [nx] - n[x] + \nu < n + 1$ and so $n = [nx] - n[x] + \nu$. But

$$\sum_{k=0}^{n-1} \left[x + \frac{k}{n} \right] = \sum_{k=0}^{\nu-1} [x] + \sum_{k=\nu}^{n-1} ([x] + 1) = \nu[x] + (n - \nu)([x] + 1) = n[x] + n - \nu,$$

and using $\nu = n - [nx] + n[x]$,

$$\sum_{k=0}^{n-1} \left[x + \frac{k}{n} \right] = n[x] + n - (n - [nx] + n[x]) = [nx].$$

This identity is proved by Hermite [65, pp. 310–315, §V].

The **Legendre symbol** is defined in the following way. Let p be an odd prime, and let a be an integer that is not a multiple of p . If there is an integer b such that $a \equiv b^2 \pmod{p}$ then $\left(\frac{a}{p}\right) = 1$, and otherwise $\left(\frac{a}{p}\right) = -1$. In other words, for an integer a that is relatively prime to p , if a is a square mod p then $\left(\frac{a}{p}\right) = 1$, and if a is not a square mod p then $\left(\frac{a}{p}\right) = -1$. For example, one checks that there is no integer b such that $b^2 \equiv 6 \pmod{7}$, and hence $\left(\frac{6}{7}\right) = -1$, while $3^2 \equiv 2 \pmod{7}$, and so $\left(\frac{2}{7}\right) = 1$.

If p and q are distinct odd primes, define integers u_k , $1 \leq k \leq \frac{p-1}{2}$, by

$$kq = p \left[\frac{kq}{p} \right] + u_k;$$

namely, u_k is the remainder of kq when divided by p . We have $1 \leq u_k \leq p-1$. Let $\mu(q, p)$ be the number of k such that $u_k > \frac{p-1}{2}$. It can be shown that [61, p. 74, Theorem 92]

$$\left(\frac{q}{p}\right) = (-1)^{\mu(q,p)};$$

this fact is called **Gauss's lemma**. For example, for $p = 13$ and $q = 3$ we work out that

$$u_1 = 3, \quad u_2 = 6, \quad u_3 = 9, \quad u_4 = 12, \quad u_5 = 2, \quad u_6 = 5,$$

and hence $\mu(3, 13) = 2$, so $(-1)^{\mu(3,13)} = 1$; on the other hand, $4^2 \equiv 3 \pmod{13}$, hence $\left(\frac{3}{13}\right) = 1$. With

$$S(q, p) = \sum_{j=1}^{\frac{p-1}{2}} \left[\frac{jq}{p} \right],$$

it is known that [61, pp. 77-78, §6.13]

$$S(q, p) \equiv \mu(q, p) \pmod{2}.$$

And it can be shown that [61, p. 76, Theorem 100]

$$S(q, p) + S(p, q) = \frac{p-1}{2} \cdot \frac{q-1}{2}. \quad (4)$$

Thus

$$\begin{aligned} \left(\frac{p}{q}\right) \left(\frac{q}{p}\right) &= (-1)^{\mu(p,q) + \mu(q,p)} \\ &= (-1)^{S(p,q) + S(q,p)} \\ &= (-1)^{\frac{p-1}{2} \cdot \frac{q-1}{2}}. \end{aligned}$$

This is Gauss's third proof of the **law of quadratic reciprocity** in the numbering [7, p. 50, §20]. This proof was published in Gauss's 1808 "Theorema-tis arithmetici demonstratio nova", which is translated in [141, pp. 112–118]. Dirichlet [38, pp. 65–72, §§42–44] gives a presentation of the proof. Eisenstein's streamlined version of Gauss's third proof is presented with historical remarks in [93]. Lemmermeyer [95] gives a comprehensive history of the law of quadratic reciprocity, and in particular writes about Gauss's third proof [95, pp. 9–10]. The formula (4) resembles the reciprocity formula for Dedekind sums [123, p. 4, Theorem 1].

Gauss obtains (4) from the following [141, p. 116, §5]: if x is irrational and n is a positive integer, then

$$\sum_{k=1}^n [kx] + \sum_{k=1}^{[nx]} \left[\frac{k}{x} \right] = n[nx], \quad (5)$$

which he proves as follows. If $\left[\frac{j}{x}\right] < k \leq \left[\frac{j+1}{x}\right]$, then $[kx] = j$. Therefore

$$\begin{aligned} \sum_{k=1}^n [kx] &= \sum_{j=1}^{\lfloor nx \rfloor} j \left(\left[\frac{j+1}{x} \right] - \left[\frac{j}{x} \right] \right) \\ &= n \lfloor nx \rfloor - \sum_{j=1}^{\lfloor nx \rfloor} \left[\frac{j}{x} \right]. \end{aligned}$$

Bachmann [6, pp. 654–658, §4] surveys later work on sums similar to (5); see also Dickson [35, Chapter X]. If m and n are relatively prime, then

$$\left\{ R\left(\frac{m}{n}\right), R\left(\frac{2m}{n}\right), \dots, R\left(\frac{(n-1)m}{n}\right) \right\} = \left\{ \frac{1}{n}, \frac{2}{n}, \dots, \frac{n-1}{n} \right\},$$

and so

$$\sum_{k=1}^{n-1} R\left(\frac{km}{n}\right) = \sum_{k=1}^{n-1} \frac{k}{n} = \frac{1}{n} \cdot \frac{(n-1)n}{2} = \frac{n-1}{2}.$$

Hence

$$\sum_{k=1}^{n-1} \left[\frac{km}{n} \right] = \sum_{k=1}^{n-1} \frac{km}{n} - \sum_{k=1}^{n-1} R\left(\frac{km}{n}\right) = \frac{(m-1)(n-1)}{2}. \quad (6)$$

There is also a simple lattice point counting argument [118, p. 113, No. 18] that gives (6).

In 1849, Dirichlet [37] shows that

$$\sum_{k=1}^n d(k) = \sum_{k=1}^n \left[\frac{n}{k} \right],$$

where $d(n)$ denotes the number of positive divisors of an integer n . (This equality is Dirichlet’s “hyperbola method”.) He then proves that

$$\sum_{k=1}^n \left[\frac{n}{k} \right] = n \log n + (2\gamma - 1)n + O(\sqrt{n}).$$

Hardy and Wright [61, pp. 264–265, Theorem 320] give a proof of this. Finding the best possible error term in the estimate for $\sum_{k=1}^n d(k)$ is “Dirichlet’s divisor problem”. Dirichlet cites the end of Section V Gauss’s *Disquisitiones Arithmeticae* as precedent for determining average magnitudes of arithmetic functions. (In Section V, Articles 302–304, of the *Disquisitiones Arithmeticae*, Gauss writes about averages of class numbers of binary quadratic forms, cf. [36, Chapter VI].)

Define (x) to be 0 if $x \in \mathbb{Z} + \frac{1}{2}$; if $x \notin \mathbb{Z} + \frac{1}{2}$ then there is an integer m_x for which $|x - m_x| < |x - n|$ for all integers $n \neq m_x$, and we define (x) to be $x - m_x$. Riemann [125, p. 105, §6] defines

$$f(x) = \sum_{n=1}^{\infty} \frac{(nx)}{n^2};$$

for any x , the series converges absolutely because $|(-nx)| < \frac{1}{2}$. Riemann states that if p and m are relatively prime and $x = \frac{p}{2m}$, then

$$f(x^+) = \lim_{h \rightarrow 0^+} f(x+h) = f(x) - \frac{\pi^2}{16m^2}, \quad f(x^-) = \lim_{h \rightarrow 0^-} f(x+h) = f(x) + \frac{\pi^2}{16m^2},$$

thus

$$f(x^-) - f(x^+) = \frac{\pi^2}{8m^2},$$

and hence that f is discontinuous at such points, and says that at all other points f is continuous; see Laugwitz [94, §2.1.1, pp. 183-184], Neuenschwander [108], and Pringsheim [119, p. 37] about Riemann's work on pathological functions. The role of pathological functions in the development of set theory is explained by Dauben [32, Chapter 1, p. 19] and Ferreirós [49, Chapter V, §1, p. 152].

For any interval $[a, b]$ and any $\sigma > 0$, it is apparent from the above that there are only finitely many $x \in [a, b]$ for which $f(x^-) - f(x^+) > \sigma$, and Riemann deduces from this that f is Riemann integrable on $[a, b]$; cf. Hawkins [63, p. 18] on the history of Riemann integration. Later in the same paper [125, p. 129, §13], Riemann states that the function

$$x \mapsto \sum_{n=1}^{\infty} \frac{(nx)}{n},$$

is not Riemann integrable in any interval.

In 1897, Cesàro [28] asks the following question (using the pseudonym, and anagram, "Rosace" [112, p. 331]). Let $\epsilon(x) = x - [x] - \frac{1}{2}$. Is the series

$$\sum_{n=1}^{\infty} \frac{\epsilon(nx)}{n} \tag{7}$$

convergent for all non-integer x ? This is plausible because the expected value of $\epsilon(x)$ is 0. Landau [89] answers this question in 1901. Landau proves that there is some g such that

$$\sum_{k=1}^n [kx] = \frac{n(n+1)x}{2} - \frac{n}{2} + O(g(n))$$

where g is a nonnegative function such $g(n) = o(n)$ and such that $\sum_{n=1}^{\infty} \frac{g(n)}{n(n+1)}$ converges, then (7) converges. And he proves that if x is rational then (7) diverges. We return to this series in §9.

Also in 1898, Franel [52] asks whether for irrational x and for $\epsilon > 0$ we have

$$\sum_{k=1}^n [kx] = \frac{n(n+1)x}{2} - \frac{n}{2} + O(n^\epsilon).$$

Then in 1899, Franel [53] asks if we can do better than this: is the error term in fact $O(1)$? Cesàro and Franel each contributed many problems to *L'Intermédiaire des mathématiciens*, the periodical in which they posed their questions. Information about Franel is given in [82].

Lerch [96] answers Franel's questions in 1904. If x is irrational and $\frac{p}{q}$ is a convergent of x (which we will define in §4), then using Theorem 2 (from §4) we can show that if $1 \leq k \leq q$ then $[kx] = \lfloor \frac{kp}{q} \rfloor$. Lerch uses this and (6) to show that if x is irrational and $\frac{p}{q}$ is a convergent of x then

$$\sum_{k=1}^q [kx] = \frac{q(q+1)x}{2} - \frac{q}{2} + R, \quad 0 < R < \frac{1}{2}.$$

Lerch states that if the continued fraction expansion of x has bounded partial quotients (defined in §4) then, for any positive integer n ,

$$\sum_{k=1}^n [kx] = \frac{n(n+1)x}{2} - \frac{n}{2} + O(\log n).$$

Lerch only gives a brief indication of the proof of this. This result is proved by Hardy and Littlewood in 1922 [59, p. 24, Theorem B3], and also in 1922 by Ostrowski [110, p. 81]. On the other hand, Lerch also constructs examples of x such that, for some positive integer c ,

$$\left| \sum_{k=1}^n [kx] - \frac{n(n+1)x}{2} + \frac{n}{2} \right| \gg n^{1-\frac{1}{c}}.$$

Nevertheless, in 1909 Sierpinski [136] proves that if x is irrational then

$$\sum_{k=1}^n [kx] = \frac{n(n+1)x}{2} - \frac{n}{2} + o(n).$$

A bibliography of Lerch's works is given in [140]. Lerch had written earlier papers on Gauss sums, Fourier series, theta functions, and the class number; many of his papers are in Czech, but some of them are in French, several of which were published in the Paris *Comptes rendus*. Several of Lerch's papers are discussed in Cresse's survey of the class number of binary quadratic forms [36, Chapter VI].

In 1899, a writer using the pseudonym "Quemquaeris" [121] ("quem quaeris" = "whom you seek") asks if we can characterize $\phi(n)$ such that for all irrational θ the series

$$\sum_{n=1}^{\infty} \frac{\phi(n)}{\sin n\pi\theta}$$

converges. In particular, the writer asks if $\phi(n) = \frac{1}{n!}$ satisfies this. In the same year, de la Vallée-Poussin [34] answers this question. (There are also responses

following de la Vallée-Poussin's by Borel and Fabry.) For a given function $\phi(n)$, de la Vallée-Poussin shows that if we have $a_n > \frac{1}{\phi(q_{n-1})}$ for all n , for a_n the n th partial quotient of θ and q_n the denominator of the n th convergent of θ , then the series

$$\sum_{n=1}^{\infty} \frac{\phi(n)}{\sin n\pi\theta}$$

will diverge. Hardy and Littlewood prove numerous results on similar series, e.g. for $\phi(n) = n^{-r}$ for real $r > 1$ and for certain classes of θ , in their papers on Diophantine approximation [58]. In 1931, Walfisz [153, p. 570, Hilfssatz 4] shows, following work of Behnke [13, p. 289, §16], that for almost all irrational $x \in [0, 1]$, if $\epsilon > 0$ then

$$\sum_{k=1}^n \frac{1}{\|k\theta\|} = O(n(\log n)^{2+\epsilon}),$$

where $\|x\| = \min(R(x), 1 - R(x))$. Walfisz's paper includes many results on related sums.

In 1916, Watson [154] finds the following asymptotic series for $S_n = \sum_{m=1}^{n-1} \csc\left(\frac{m\pi}{n}\right)$:

$$\pi S_n \sim 2n \log(2n) + 2n(\gamma - \log \pi) + \sum_{j=1}^{\infty} (-1)^j \frac{B_{2j}^2 (2^{2j} - 2) \pi^{2j}}{j(2j)! n^{2j-1}},$$

where γ is Euler's constant and B_j are the Bernoulli numbers. Truncating the asymptotic series and rewriting gives

$$S_n = \frac{2n \log n}{\pi} + n \cdot \frac{2 \log 2 + 2\gamma - 2 \log \pi}{\pi} + O\left(\frac{1}{n}\right).$$

For example, computing S_{1000} directly we get $S_{1000} = 4477.593932\dots$, and computing the right-hand side of the above formula without the error term we obtain $4477.594019\dots$. A cleaner derivation of the asymptotic series using the Euler-Maclaurin summation formula is given later by Williams [156].

Early surveys of Diophantine approximation are given by Bohr and Cramér [20, pp. 833–836, §39] and Koksma [81, pp. 102–110]. Hlawka and Binder [66] present the history of the initial years of the theory of uniform distribution. Narkiewicz [106, pp. 82–95, §2.5 and pp. 175–183 §3.5] gives additional historical references on Diophantine approximation. The papers of Hardy and Littlewood on Diophantine approximation are reprinted in [58]. Perron [113] and Brezinski [23] give historical references on continued fractions, and there is reliable material on the use of continued fractions by 17th century mathematicians in Whiteside [155]. Fowler [51] presents a prehistory of continued fractions in Greek mathematics.

4 Preliminaries on continued fractions

Let

$$\|x\| = \min_{k \in \mathbb{Z}} |x - k| = \min(R(x), 1 - R(x)).$$

Let μ be Lebesgue measure on $[0, 1]$, and let $\Omega = [0, 1] \setminus \mathbb{Q}$.
 For positive integers a_1, \dots, a_n , we define

$$[a_1, \dots, a_n] = \frac{1}{a_1 + \frac{1}{\dots + \frac{1}{a_{n-1} + \frac{1}{a_n}}}}.$$

For example, $[1, 1, 1] = \frac{2}{3}$.

Let \mathbb{N} be the set of positive integers. We call $a \in \mathbb{N}^{\mathbb{N}}$ a **continued fraction**, and we call a_n the **n th partial quotient** of a . If there is some $K > 0$ such that $a_n \leq K$ for all n then we say that a has bounded partial quotients. We call $[a_1, \dots, a_n]$ the **n th convergent** of a . For $n \geq 1$ let

$$\frac{p_n}{q_n} = [a_1, \dots, a_n],$$

with p_n, q_n positive integers that are relatively prime, and set

$$p_0 = 0, \quad q_0 = 1.$$

One can show by induction [44, p. 70, Lemma 3.1] that for $n \geq 1$ we have

$$\begin{pmatrix} p_n & p_{n-1} \\ q_n & q_{n-1} \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} a_1 & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} a_n & 1 \\ 1 & 0 \end{pmatrix}. \quad (8)$$

We have

$$p_1 = 1, \quad q_1 = a_1,$$

and from (8) we get for all $n \geq 1$ that

$$p_{n+1} = a_{n+1}p_n + p_{n-1},$$

and

$$q_{n+1} = a_{n+1}q_n + q_{n-1}.$$

Since the a_n are positive integers we get by induction that for all $n \geq 1$,

$$p_n \geq 2^{(n-1)/2}, \quad q_n \geq 2^{(n-1)/2}. \quad (9)$$

In fact, setting $F_1 = 1, F_2 = 1, F_{n+1} = F_n + F_{n-1}$ for $n \geq 2$, with F_n the **n th Fibonacci number**, as $a_n \geq 1$ we check by induction that

$$p_n \geq F_n, \quad q_n \geq F_{n+1}.$$

Taking determinants of (8) gives us for all $n \geq 1$ that

$$p_n q_{n-1} - p_{n-1} q_n = (-1)^{n+1}, \quad (10)$$

and then by induction we have for all $n \geq 1$,

$$\frac{p_n}{q_n} = \sum_{k=1}^n \frac{(-1)^{k+1}}{q_{k-1}q_k}.$$

For any $a \in \mathbb{N}^{\mathbb{N}}$, as $n \rightarrow \infty$ this sequence of sums converges and we denote its limit by $v(a)$. We have for all $n \geq 1$,

$$v(a) - \frac{p_n}{q_n} = \sum_{k=n+1}^{\infty} \frac{(-1)^{k+1}}{q_{k-1}q_k}.$$

Since the right hand side is an alternating series we obtain for $n \geq 1$,

$$\left| v(a) - \frac{p_n}{q_n} \right| < \frac{1}{q_n q_{n+1}}, \quad (11)$$

and

$$\left| v(a) - \frac{p_n}{q_n} \right| > \frac{1}{q_n q_{n+1}} - \frac{1}{q_{n+1} q_{n+2}} = \frac{a_{n+2}}{q_n q_{n+2}} \geq \frac{1}{q_n (q_{n+1} + q_n)}, \quad (12)$$

and

$$\frac{p_2}{q_2} < \frac{p_4}{q_4} < \dots < v(a) < \dots < \frac{p_3}{q_3} < \frac{p_1}{q_1}. \quad (13)$$

For $x \in \Omega$, we say that $\frac{p}{q} \in \mathbb{Q}$, $q > 0$, is a **best approximation to x** if $\|qx\| = |qx - p|$ and $\|q'x\| > \|qx\|$ for $1 \leq q' < q$. The following theorem shows in particular that the convergents of a continued fraction a are best approximations to $v(a)$ [127, p. 22, Chapter 2, §3, Theorem 1].

Theorem 2 (Best approximations). *Let $a \in \mathbb{N}^{\mathbb{N}}$. For any $n \geq 1$,*

$$\|q_n v(a)\| = |p_n - q_n v(a)|.$$

If $1 \leq q < q_{n+1}$, then for any $p \in \mathbb{Z}$,

$$|qv(a) - p| \geq |p_n - q_n v(a)|.$$

Proof. By (11), $|q_n v(a) - p_n| < \frac{1}{q_{n+1}}$. But

$$q_{n+1} \geq q_2 = a_2 q_1 + q_0 \geq q_1 + q_0 = a_1 + 1 \geq 2.$$

So $|q_n v(a) - p_n| < \frac{1}{2}$, which means $\|q_n v(a)\| = |q_n v(a) - p_n|$.

Write $x = v(a)$ and let $A = \begin{pmatrix} p_{n+1} & p_n \\ q_{n+1} & q_n \end{pmatrix}$. Applying (10),

$$\det A = p_{n+1} q_n - p_n q_{n+1} = (-1)^n.$$

Let

$$\begin{pmatrix} \mu \\ \nu \end{pmatrix} = A^{-1} \begin{pmatrix} p \\ q \end{pmatrix} = \frac{1}{\det A} \begin{pmatrix} q_n & -p_n \\ -q_{n+1} & p_{n+1} \end{pmatrix} \begin{pmatrix} p \\ q \end{pmatrix} = (-1)^n \begin{pmatrix} q_n & -p_n \\ -q_{n+1} & p_{n+1} \end{pmatrix} \begin{pmatrix} p \\ q \end{pmatrix}.$$

Then

$$qx - p = (\mu q_{n+1} + \nu q_n)x - (p_{n+1}\mu + p_n\nu) = \mu(q_{n+1}x - p_{n+1}) + \nu(q_nx - p_n)$$

and

$$\begin{pmatrix} p \\ q \end{pmatrix} = A \begin{pmatrix} \mu \\ \nu \end{pmatrix} = \begin{pmatrix} p_{n+1} & p_n \\ q_{n+1} & q_n \end{pmatrix} \begin{pmatrix} \mu \\ \nu \end{pmatrix} = \begin{pmatrix} p_{n+1}\mu + p_n\nu \\ q_{n+1}\mu + q_n\nu \end{pmatrix}.$$

In particular, $\mu, \nu \in \mathbb{Z}$. Suppose by contradiction that $\nu = 0$. Then $(q - \mu q_{n+1})x = p - \mu p_{n+1}$, and as $x \notin \mathbb{Q}$ it must then be that $q = \mu q_{n+1}$ and $p = \mu p_{n+1}$. But $q = \mu q_{n+1}$, $1 \leq q < q_{n+1}$, and $\mu \in \mathbb{Z}$ are together a contradiction. Therefore $\nu \neq 0$. Either $\mu = 0$ or $\mu \neq 0$. For $\mu = 0$,

$$|qx - p| = |\nu||q_nx - p_n| \geq |q_nx - p_n|,$$

which is the claim. For $\mu \neq 0$ we use the fact $q = q_{n+1}\mu + q_n\nu$ and $1 \leq q < q_{n+1}$. If $\mu, \nu > 0$ then $q < q_{n+1}$ is contradicted, and if $\mu, \nu < 0$ then $q \geq 1$ is contradicted. Therefore μ and ν have different signs, say $\mu = (-1)^N|\mu|$ and $\nu = (-1)^{N+1}|\nu|$. Furthermore, we get from (13) that

$$\operatorname{sgn}(q_nx - p_n) = (-1)^n, \quad \operatorname{sgn}(q_{n+1}x - p_{n+1}) = (-1)^{n+1}.$$

Therefore

$$\begin{aligned} qx - p &= \mu(q_{n+1}x - p_{n+1}) + \nu(q_nx - p_n) \\ &= (-1)^N|\mu| \cdot (-1)^{n+1}|q_{n+1}x - p_{n+1}| + (-1)^{N+1}|\nu| \cdot (-1)^n|q_nx - p_n|, \end{aligned}$$

hence

$$|qx - p| = |\mu||q_{n+1}x - p_{n+1}| + |\nu||q_nx - p_n| \geq |\nu||q_nx - p_n| \geq |q_nx - p_n|,$$

which is the claim. \square

The above theorem says, a fortiori, that the convergents of a are best approximations to $v(a)$. It can also be proved that if $\frac{p}{q} \in \mathbb{Q}$, $q > 0$, is a best approximation to $v(a)$ then $\frac{p}{q}$ is a convergent of a [92, p. 9, Theorem 6]. Cassels [27, p. 2, Chapter I] works out the theory of continued fractions according to this point of view. Similarly, Milnor [101, p. 234, Appendix C] works out the theory of continued fractions in the language of rotations of the unit circle.

We define the **Gauss transformation** $T : \Omega \rightarrow \Omega$ by $T(x) = R(\frac{1}{x})$ for $x \in \Omega$, and we define $\Phi : \Omega \rightarrow \mathbb{N}^{\mathbb{N}}$ by

$$(\Phi(x))_n = \left\lceil \frac{1}{T^{n-1}(x)} \right\rceil, \quad n \geq 1.$$

One can check that if $a \in \mathbb{N}^{\mathbb{N}}$ then $v(a) \in \Omega$ [44, p. 73, Lemma 3.2]. (Namely, the value of a nonterminating continued fraction is irrational.) One can prove that $v : \mathbb{N}^{\mathbb{N}} \rightarrow \Omega$ is injective [44, p. 75, Lemma 3.4], and for $x \in \Omega$ that [44, p. 78, Lemma 3.6]

$$(v \circ \Phi)(x) = x.$$

Therefore $\Phi : \Omega \rightarrow \mathbb{N}^{\mathbb{N}}$ is a bijection. Moreover, Φ is a homeomorphism, when \mathbb{N} has discrete topology, $\mathbb{N}^{\mathbb{N}}$ has the product topology, and Ω has the subspace topology inherited from \mathbb{R} [44, p. 86, Exercise 3.2.2]. That $\mathbb{N}^{\mathbb{N}}$ and Ω are homeomorphic can also be proved without using continued fractions [2, p. 106, Theorem 3.68]. In descriptive set theory, the topological space $\mathcal{N} = \mathbb{N}^{\mathbb{N}}$ is called the **Baire space**, and the Alexandrov-Urysohn theorem states that \mathcal{N} has the universal property that any nonempty Polish space that is zero-dimensional (there is a basis of clopen sets for the topology) and all of whose compact subsets have empty interior is homeomorphic to \mathcal{N} [75, p. 37, Theorem 7.7]. Some of Baire's work on \mathcal{N} is described in [71, pp. 119–120] and [5, pp. 349, 372].

For $I = [0, 1]$ and for $T : I \rightarrow I$, $T(x) = R(1/x)$ for $x > 0$ and $T(0) = 0$. For $k \geq 1$ let $I_k = \left(\frac{1}{k+1}, \frac{1}{k}\right)$, so if $x \in I_k$ then $T(x) = x^{-1} - k$. Then for $x \in I = [0, 1]$,

$$T(x) = \sum_{k=1}^{\infty} 1_{I_k}(x)(x^{-1} - k).$$

For $S = \{0\} \cup \{k^{-1} : k \geq 1\}$, $I \setminus S = \bigcup_{k \geq 1} I_k$, and for $x \in I \setminus S$,

$$T'(x) = - \sum_{k=1}^{\infty} 1_{I_k}(x)x^{-2},$$

and for $x \in I_k$, $k^2 < |T'(x)| < (k+1)^2$. Differentiability and dynamical properties of the Gauss transformation are worked out by Cornfeld, Fomin and Sinai [30, pp. 165–177, Chapter 7, §4], as an instance of **piecewise monotonic** transformations.

For each $n \geq 1$ we define $a_n : \Omega \rightarrow \mathbb{N}$ by $a_n(x) = (\Phi(x))_n$. For example, $e - 2 \in \Omega$, and it is known [92, p. 74, Theorem 2] that for $k \geq 1$,

$$a_{3k}(e - 2) = a_{3k-2}(e - 2) = 1 \quad \text{and} \quad a_{3k-1}(e - 2) = 2k.$$

The pattern for the continued fraction expansion of e seems first to have been worked out by Roger Cotes in 1714 [50], and was later proved by Euler using a method involving the Riccati equation [31].

For $n \geq 1$ and $i \in \mathbb{N}^n$, let

$$I_n(i) = \{\omega \in \Omega : a_k(x) = i_k, 1 \leq k \leq n\}.$$

For $x \in I_n(i)$,

$$\frac{p_n(x)}{q_n(x)} = [i_1, \dots, i_n], \quad \frac{p_{n-1}(x)}{q_{n-1}(x)} = [i_1, \dots, i_{n-1}].$$

The following is an expression for the sets $I_n(i)$ [69, p. 18, Theorem 1.2.2].

Theorem 3. Let $n \geq 1$, $i \in \mathbb{Z}_{\geq 1}^n$, and for $p_n = p_n(x)$, $q_n = q_n(x)$, $x \in I_n(i)$,

$$u_n(i) = \begin{cases} \frac{p_n + p_{n-1}}{q_n + q_{n-1}} & n \text{ odd} \\ \frac{p_n}{q_n} & n \text{ even} \end{cases}$$

and

$$v_n(i) = \begin{cases} \frac{p_n}{q_n} & n \text{ odd} \\ \frac{p_n + p_{n-1}}{q_n + q_{n-1}} & n \text{ even}. \end{cases}$$

Then

$$I_n(i) = \Omega \cap (u_n(i), v_n(i)).$$

It follows from the above that for $i \in \mathbb{N}^n$, $n \geq 1$,

$$\mu(I_n(i)) = \frac{1}{q_n(q_n + q_{n-1})}.$$

5 Diophantine conditions

For real $\tau, \gamma > 0$ let

$$\begin{aligned} \mathcal{D}(\tau, \gamma) &= \bigcap_{q \in \mathbb{Z}_{\geq 1}, p \in \mathbb{Z}} \left\{ x \in [0, 1] : \left| x - \frac{p}{q} \right| \geq \gamma q^{-\tau} \right\} \\ &= \bigcap_{q \in \mathbb{Z}_{\geq 1}} \left\{ x \in [0, 1] : \|qx\| \geq \gamma q^{-\tau+1} \right\}, \end{aligned}$$

and let

$$\mathcal{D}(\tau) = \bigcup_{\gamma > 0} \mathcal{D}(\tau, \gamma).$$

We relate the sets $\mathcal{D}(\tau)$ and continued fractions expansions [101, p. 241, Lemma C.6], cf. [159, p. 130, Proposition 2.4].

Lemma 4. For $\tau > 0$ and $x \in \Omega$, $x \in \mathcal{D}(\tau)$ if and only if there is some $C = C(x) > 0$ such that $q_{n+1}(x) \leq Cq_n(x)^{\tau-1}$ for all $n \geq 1$.

Proof. For $x \in \Omega$, write $q_n = q_n(x)$. By (12), $\|q_n x\| > \frac{1}{q_{n+1} + q_n}$, and by (11), $\|q_n x\| < \frac{1}{q_{n+1}}$. Suppose $x \in \mathcal{D}(\tau)$, so there is some $\gamma > 0$ such that $x \in \mathcal{D}(\tau, \gamma)$. Then

$$q_{n+1} < \frac{1}{\|q_n x\|} \leq \gamma^{-1} q_n^{\tau-1}.$$

Suppose $q_{n+1} \leq Cq_n^{\tau-1}$ for all $n \geq 1$. For $q \in \mathbb{Z}_{\geq 1}$, take $q_n \leq q < q_{n+1}$. Using Theorem 2,

$$\|qx\| \geq \|q_n x\| > \frac{1}{q_{n+1} + q_n} > \frac{1}{2q_{n+1}} \geq \frac{1}{2} C^{-1} q_n^{-\tau+1} \geq \frac{1}{2C} \cdot q^{-\tau+1},$$

which means that $x \in \mathcal{D}(\tau, \frac{1}{2C})$. \square

For K a positive integer, let

$$\mathcal{B}_K = \{x \in \Omega : a_n(x) \leq K \text{ for all } n \geq 1\},$$

so $\mathcal{B} = \bigcup_{K \geq 1} \mathcal{B}_K$ is the set of those $x \in \Omega$ with bounded partial quotients.

Lemma 5. *For $x \in \Omega$, $x \in \mathcal{D}(2)$ if and only if $x \in \mathcal{B}$.*

Proof. Write $a_n = a_n(x)$ and $q_n = q_n(x)$. If $x \in \mathcal{D}(2)$ then there is some $\gamma > 0$ such that $x \in \mathcal{D}(2, \gamma)$, hence for $n \geq 1$,

$$q_{n+1} < \frac{1}{\|q_n x\|} \leq \gamma^{-1} q_n.$$

Now, $q_{n+1} = a_{n+1}q_n + q_{n-1}$ for $n \geq 1$, so

$$a_{n+1} < q_{n+1}q_n^{-1} < q_n^{-1} \cdot \gamma^{-1} q_n = \gamma^{-1},$$

which shows that x has bounded partial quotients.

If $x \in \mathcal{B}_K$, let $q \in \mathbb{Z}_{\geq 1}$ and let $q_n \leq q < q_{n+1}$. Using Theorem 2 and then (12),

$$\|qx\| \geq \|q_n x\| > \frac{1}{q_{n+1} + q_n} = \frac{1}{a_{n+1}q_n + q_{n-1} + q_n} > \frac{1}{(a_{n+1} + 2)q_n}.$$

As $x \in \mathcal{B}_K$,

$$\|qx\| > \frac{1}{(K+2)q_n} \geq \frac{1}{K+2} q^{-1},$$

which means that $x \in \mathcal{D}(2, \frac{1}{K+2})$. \square

A complex number α is called an **algebraic number of degree d** , $d \geq 0$, if there is some polynomial $f \in \mathbb{Z}[x]$ with degree d such that $f(\alpha) = 0$ and if $g \in \mathbb{Z}[x]$ has degree $< d$ and $g(\alpha) = 0$ then $g = 0$. An algebraic number of degree 2 is called a **quadratic irrational**. Let $\alpha \in \Omega$. It was proved by Euler [61, p. 144, Theorem 176] that if there is some $p > 0$ and some L such that $a_{l+p}(\alpha) = a_l(\alpha)$ for all $l \geq L$ then α is a quadratic irrational. The converse of this was proved by Lagrange [61, p. 144, Theorem 177], namely that a quadratic irrational has eventually periodic partial quotients. For example, $\alpha = \sqrt{11} - 3 \in \Omega$ is a quadratic irrational, being a root of $x^2 + 6x - 2$, and one works out that $a_1(\alpha) = 3, a_2(\alpha) = 6$, and that $a_{l+2}(\alpha) = a_l(\alpha)$ for $l \geq 1$. In particular, if $\alpha \in \Omega$ is a quadratic irrational, then α has bounded partial quotients.

Liouville [61, p. 161, Theorem 191] proved that if $x \in \Omega$ is an algebraic number of degree $d \geq 2$, then $x \in \mathcal{D}(d)$. The **Thue-Siegel-Roth theorem** [48, p. 55, Theorem 1.23] states that if $x \in \Omega$ is an algebraic number, then for any $\delta > 0$ there is some $q_\delta \in \mathbb{Z}_{\geq 1}$ such that for all $q \geq q_\delta$,

$$\|qx\| \geq q^{-1-\delta}.$$

See Schmidt [132, p. 195, Theorem 2B].

6 Sums of reciprocals

We are interested in getting bounds on the sum $\sum_{j=1}^m \frac{1}{\|jx\|}$. This is an appealing question because the terms $\frac{1}{\|jx\|}$ are unbounded.

Rather than merely stating that $\sum_{k=1}^{\infty} \frac{1}{k} = \infty$, we give more information by giving the estimate

$$\sum_{k=1}^n \frac{1}{k} = \log n + \gamma + O(n^{-1}),$$

where γ is Euler's constant. Likewise, rather than merely stating that there are infinitely many primes, we state more information with [90, p. 102, §28]

$$\sum_{p \leq x} \frac{1}{p} = \log \log x + B + O\left(\frac{1}{\log x}\right),$$

for a certain constant B (namely "Merten's constant"), or with [90, p. 226, §61]

$$\sum_{p \leq x} p = \frac{x^2}{2 \log x} + O\left(\frac{x^2}{(\log x)^2}\right).$$

Because $|\sin(\pi x)| = \sin(\pi \|x\|) \leq \pi \|x\|$ and

$$|\sin(\pi x)| = \sin(\pi \|x\|) \geq \frac{2}{\pi} \pi \|x\| = 2 \|x\|,$$

we have

$$\frac{1}{\pi} \sum_{j=1}^m \frac{1}{\|jx\|} \leq \sum_{j=1}^m \frac{1}{|\sin \pi jx|} \leq \frac{1}{2} \sum_{j=1}^m \frac{1}{\|jx\|}. \quad (14)$$

Thus, getting bounds on $\sum_{j=1}^m \frac{1}{\|jx\|}$ will give us bounds on $\sum_{j=1}^m \frac{1}{|\sin \pi jx|}$.

Let ψ be a nondecreasing function defined on the positive integers such that $\psi(h) > 0$ for $h \geq 1$ (for example, $\psi(h) = \log(2h)$). Following Kuipers and Niederreiter [85, p. 121, Definition 3.3], we say that an irrational number x is **of type** $< \psi$ if $\|hx\| \geq \frac{1}{h\psi(h)}$ for all integers $h \geq 1$. If ψ is a constant function, then we say that x is **of constant type**.

Lemma 6. $x \in \Omega$ is of constant type if and only if it has bounded partial quotients.

Proof. Suppose that $x \in \Omega$ is of constant type. So there is some $K > 0$ such that $\|hx\| \geq \frac{1}{hK}$ for all integers $h \geq 1$. For $n \geq 2$ we have $q_n = a_n q_{n-1} + q_{n-2}$, and hence, by (11),

$$a_n = \frac{q_n}{q_{n-1}} - \frac{q_{n-2}}{q_{n-1}} < \frac{q_n}{q_{n-1}} \leq q_n \cdot K \|q_{n-1}x\| < K,$$

showing that x has bounded partial quotients.

Suppose that $x \in \Omega$ has bounded partial quotients, say $a_n \leq K$ for all $n \geq 1$. Let h be a positive integer and take $q_n \leq h < q_{n+1}$. Then first by Theorem 2 and then by (12),

$$\|hx\| \geq \|q_n x\| > \frac{1}{q_n + q_{n+1}} > \frac{1}{2q_{n+1}} = \frac{1}{2(a_{n+1}q_n + q_{n-1})} > \frac{1}{2(a_{n+1}q_n + q_n)},$$

and so

$$\|hx\| > \frac{1}{2(K+1)} \frac{1}{q_n} \geq \frac{1}{2(K+1)} \frac{1}{h},$$

showing that x is of constant type. \square

However, almost all x do not have bounded partial quotients [78, p. 60, Theorem 29]. Shallit [135] gives a survey on numbers with bounded partial quotients.

We state and prove a result of Khinchin's [78, p. 69, Theorem 32] that we then use.

Lemma 7. *Let f be a positive function on the positive integers. If*

$$\sum_{q=1}^{\infty} f(q) < \infty,$$

then for almost all $x \in \Omega$ there are only finitely many q such that $\|qx\| < f(q)$.

Proof. For each positive integer q , let $E_q = \{t \in \Omega : \|qt\| < f(q)\}$. If $t \in E_q$, then there is some integer p with $0 \leq p \leq q$ such that $\left|t - \frac{p}{q}\right| < \frac{f(q)}{q}$. It follows that

$$E_q \subset \left(0, \frac{f(q)}{q}\right) \cup \left(1 - \frac{f(q)}{q}, 1\right) \cup \bigcup_{p=1}^{q-1} \left(\frac{p}{q} - \frac{f(q)}{q}, \frac{p}{q} + \frac{f(q)}{q}\right).$$

Therefore

$$\sum_{q=1}^{\infty} \mu(E_q) \leq \sum_{q=1}^{\infty} 2q \cdot \frac{f(q)}{q} = 2 \sum_{q=1}^{\infty} f(q) < \infty.$$

Let $E = \limsup_{q \rightarrow \infty} E_q$, i.e. $E = \{t \in \Omega : t \in E_q \text{ for infinitely many } q\}$. Then by the Borel-Cantelli lemma [19, p. 59, Theorem 4.3] we have that $\mu(E) = 0$. Therefore, for almost all $t \in \Omega$ there are only finitely many q such that $t \in E_q$, i.e., for almost all $t \in \Omega$ there are only finitely many q such that $\|qt\| < f(q)$. \square

The above lemma is proved in Benedetto and Czaja [15, p. 183, Theorem 4.3.3] using the fact that a function of bounded variation is differentiable almost everywhere. We outline the proof. Define $F : [0, 1] \rightarrow \mathbb{R}_{>0}$ by $F(x) = \frac{f(q)}{q}$ if

$x = \frac{a}{q}$, $\gcd(a, q) = 1$, $0 \leq a \leq q$, and $F(x) = 0$ if $x \in \Omega$. Writing $\mathcal{A}_q = \left\{ \frac{a}{q} : 0 \leq a \leq q, \gcd(a, q) = 1 \right\}$, for $0 = t_0 < t_1 < \dots < t_N = 1$,

$$\sum_{j=0}^N F(t_j) = \sum_{q=1}^{\infty} \sum_{j=0}^N F(t_j) \cdot 1_{\mathcal{A}_q}(t_j) = \sum_{q=1}^{\infty} \frac{f(q)}{q} \sum_{j=0}^N 1_{\mathcal{A}_q}(t_j) \leq \sum_{q=1}^{\infty} f(q).$$

It follows that the total variation of F is $\leq 2 \sum_{q=1}^{\infty} f(q)$ and hence F is a function of **bounded variation**. Because F has bounded variation, the set D_F of $x \in [0, 1]$ at which F is differentiable is a Borel set with $\lambda(D_F) = 1$. Check that $F'(x) = 0$ for $x \in D_F \setminus \mathbb{Q}$, and using this, if $\frac{a_n}{q_n} \rightarrow x$ with $\gcd(a_n, q_n) = 1$ and $0 \leq a_n \leq q_n$ then for some N , if $n \geq N$ then $\left| x - \frac{a_n}{q_n} \right| \geq \frac{F(q_n)}{q_n}$.

We use the above lemma to prove the following result.

Lemma 8. *Let $\epsilon > 0$. For almost all $x \in \Omega$, there is some $K > 0$ such that x is of type $< K(\log h)^{1+\epsilon}$.*

Proof. Let

$$E = \left\{ t \in \Omega : \|ht\| < \frac{1}{h(\log h)^{1+\epsilon}} \text{ for infinitely many } h \right\}.$$

Since $\sum_{h=1}^{\infty} \frac{1}{h(\log h)^{1+\epsilon}}$ converges, we have by Lemma 7 that $\mu(E) = 0$. Let $t \in \Omega \setminus E$. Then $\|ht\| \geq \frac{1}{h(\log h)^{1+\epsilon}}$ for all sufficiently large h . It follows that there is some K such that t is of type $< K(\log h)^{1+\epsilon}$. \square

The following technical lemma is from Kuipers and Niederreiter [85, p. 130, Exercise 3.9]; cf. Lang [92, p. 39, Lemma].

Lemma 9. *Let $x \in \Omega$ be of type $< \psi$. If $n \geq 0$ and if $0 \leq h_0 < q_{n+1}$, then*

$$\sum_{\substack{1 \leq j \leq q_n \\ j+h_0 < q_{n+1}}} \frac{1}{\|(j+h_0)x\|} < 6q_n(\psi(q_n) + \log q_n).$$

Proof. Since p_n and q_n are relatively prime, the remainders of jp_n , $j = 1, \dots, q_n$, when divided by q_n are all distinct. Then also, the remainders of $jp_n + h_0p_n$, $j = 1, \dots, q_n$, when divided by q_n are all distinct. Let λ_j , $j = 1, \dots, q_n$, be the remainder of $jp_n + h_0p_n$ when divided by q_n . We have $\{\lambda_j : 1 \leq j \leq q_n\} = \{0, \dots, q_n - 1\}$; let $\lambda_{j_1} = 0$, $\lambda_{j_2} = 1$, and $\lambda_{j_3} = q_n - 1$.

Write $x = \frac{p_n}{q_n} + \frac{\delta_n}{q_n q_{n+1}}$; by (11) we have $|\delta_n| < 1$. If $j + h_0 < q_{n+1}$ then by Theorem 2 we have $\|(j + h_0)x\| \geq \|q_n x\|$, and since x is of type $< \psi$ we have

$$\|(j + h_0)x\| \geq \|q_n x\| \geq \frac{1}{q_n \psi(q_n)}.$$

Let, for $i = 1, 2, 3$,

$$A_i = \begin{cases} 1, & \text{if } j_i + h_0 < q_{n+1}, \\ 0, & \text{if } j_i + h_0 \geq q_{n+1}. \end{cases}$$

If $j + h_0 < q_{n+1}$ and $j \neq j_1, j_2, j_3$, then

$$\left\| \frac{\lambda_j}{q_n} + \frac{(j+h_0)\delta_n}{q_n q_{n+1}} \right\| \geq \min \left\{ \left\| \frac{\lambda_j}{q_n} + \frac{1}{q_n} \right\|, \left\| \frac{\lambda_j}{q_n} - \frac{1}{q_n} \right\| \right\}.$$

It follows that

$$\begin{aligned} \sum_{\substack{1 \leq j \leq q_n \\ j+h_0 < q_{n+1}}} \frac{1}{\|(j+h_0)x\|} &= \frac{A_1}{\|(j_1+h_0)x\|} + \frac{A_2}{\|(j_2+h_0)x\|} + \frac{A_3}{\|(j_3+h_0)x\|} \\ &+ \sum_{\substack{1 \leq j \leq q_n \\ j+h_0 < q_{n+1} \\ j \neq j_1, j_2, j_3}} \frac{1}{\left\| \frac{\lambda_j}{q_n} + \frac{(j+h_0)\delta_n}{q_n q_{n+1}} \right\|} \\ &\leq 3q_n \psi(q_n) + \sum_{\substack{1 \leq j \leq q_n \\ j+h_0 < q_{n+1} \\ j \neq j_1, j_2, j_3}} \frac{1}{\left\| \frac{\lambda_j}{q_n} + \frac{(j+h_0)\delta_n}{q_n q_{n+1}} \right\|} \\ &\leq 3q_n \psi(q_n) + \sum_{\substack{1 \leq j \leq q_n \\ j+h_0 < q_{n+1} \\ j \neq j_1, j_2, j_3}} \frac{1}{\left\| \frac{\lambda_j}{q_n} + \frac{1}{q_n} \right\|} \\ &+ \sum_{\substack{1 \leq j \leq q_n \\ j+h_0 < q_{n+1} \\ j \neq j_1, j_2, j_3}} \frac{1}{\left\| \frac{\lambda_j}{q_n} - \frac{1}{q_n} \right\|} \\ &< 3q_n \psi(q_n) + 2 \sum_{k=1}^{q_n-1} \frac{1}{\left\| \frac{k}{q_n} \right\|}. \end{aligned}$$

But $\frac{1}{\|y\|} < \frac{1}{R(y)} + \frac{1}{1-R(y)}$ for $y \notin \mathbb{Z}$, so

$$\begin{aligned} \sum_{k=1}^{q_n-1} \frac{1}{\left\| \frac{k}{q_n} \right\|} &< \sum_{k=1}^{q_n-1} \frac{1}{R\left(\frac{k}{q_n}\right)} + \frac{1}{1-R\left(\frac{k}{q_n}\right)} \\ &= \sum_{k=1}^{q_n-1} \frac{1}{\frac{k}{q_n}} + \frac{1}{1-\frac{k}{q_n}} \\ &= 2q_n \sum_{k=1}^{q_n-1} \frac{1}{k} \\ &< 3q_n \log q_n; \end{aligned}$$

the last inequality is because, for all $m \geq 1$,

$$\sum_{k=1}^m \frac{1}{k} < \frac{3}{2} \log(m+1).$$

□

We now use Lemma 9 to obtain a bound on $\sum_{j=1}^m \frac{1}{\|jx\|}$ in terms of the type of x . This is from Kuipers and Niederreiter [85, p. 131, Exercise 3.11]; cf. Lang [92, p. 39, Theorem 2].

Theorem 10. *If $x \in \Omega$ is of type $< \psi$, then for all $m \geq 1$ we have*

$$\sum_{j=1}^m \frac{1}{\|jx\|} < 12m(\psi(m) + \log m).$$

Proof. We shall prove the claim by induction. Because x is of type $< \psi$, we have

$$\frac{1}{\|x\|} \leq \psi(1) < 12\psi(1),$$

so the claim is true for $m = 1$. Take $m > 1$ and assume that the claim is true for all $1 \leq m' < m$. We shall show that it is true for m .

Let $q_n \leq m < q_{n+1}$. Either $m < 2q_n$ or $m \geq 2q_n$. In the first case, using Lemma 9 we have

$$\begin{aligned} \sum_{j=1}^m \frac{1}{\|jx\|} &= \sum_{j=1}^{q_n} \frac{1}{\|jx\|} + \sum_{j=1}^{m-q_n} \frac{1}{\|(j+q_n)x\|} \\ &< 12q_n(\psi(q_n) + \log q_n) \\ &< 12m(\psi(m) + \log m). \end{aligned}$$

In the second case, using the induction assumption (with $m' = m - q_n$) and Lemma 9 we have, because $q_n < m - q_n$,

$$\begin{aligned} \sum_{j=1}^m \frac{1}{\|jx\|} &= \sum_{j=1}^{m-q_n} \frac{1}{\|jx\|} + \sum_{j=1}^{q_n} \frac{1}{\|(j+m-q_n)x\|} \\ &< 12(m-q_n)(\psi(m-q_n) + \log(m-q_n)) + 6q_n(\psi(q_n) + \log q_n) \\ &< 12(m-q_n)(\psi(m) + \log m) + 12q_n(\psi(q_n) + \log q_n) \\ &= 12m(\psi(m) + \log m) - 12q_n(\psi(m) - \psi(q_n) + \log m - \log q_n) \\ &\leq 12m(\psi(m) + \log m). \end{aligned}$$

The claim is true in both cases, which completes the proof by induction. □

We can now establish for almost all $x \in \Omega$ a tractable upper bound on the sum $\sum_{j=1}^m \frac{1}{\|jx\|}$, and thus by (14) also on $\sum_{j=1}^m \frac{1}{|\sin \pi jx|}$; cf. Lang [92, p. 44, Theorem 3].

Theorem 11. *Let $\epsilon > 0$. For almost all $x \in \Omega$, we have*

$$\sum_{j=1}^m \frac{1}{\|jx\|} = O\left(m(\log m)^{1+\epsilon}\right),$$

while if x has bounded partial quotients then

$$\sum_{j=1}^m \frac{1}{\|jx\|} = O(m \log m).$$

Proof. Let $\epsilon > 0$. By Lemma 8, for almost all $x \in \Omega$ there is some K such that x is of type $< K(\log h)^{1+\epsilon}$. For such an x , it follows from Theorem 10 that for all $m \geq 1$,

$$\sum_{j=1}^m \frac{1}{\|jx\|} < 12m(K(\log m)^{1+\epsilon} + \log m) = O\left(m(\log m)^{1+\epsilon}\right).$$

If x has bounded partial quotients then by Lemma 6 it is of constant type, say $\psi(m) = K$ for some K . It follows from Theorem 10 that for all $m \geq 1$,

$$\sum_{j=1}^m \frac{1}{\|jx\|} < 12m(K + \log m) = O(m \log m).$$

□

For example, take $x = \frac{-1+\sqrt{5}}{2}$, for which $a_n(x) = 1$ for all $n \in \mathbb{N}$, and so in particular x has bounded partial quotients. In Figure 1 we plot $\frac{1}{m \log m} \cdot \sum_{j=1}^m \frac{1}{\|jx\|}$ for $m = 5000, \dots, 40000$. These computations suggest that there is some constant C for which $\sum_{j=1}^m \frac{1}{\|jx\|} > Cm \log m$ for all m . In Theorem 13 we shall prove that for almost all $x \in \Omega$ there is such a $C(x)$.

However, the estimate in Theorem 11 is not true for all $x \in \Omega$. Define $a \in \mathbb{N}^{\mathbb{N}}$ as follows. Let a_1 be any element of \mathbb{N} . Then inductively, define a_{n+1} to be any element of \mathbb{N} that is $> q_n^{n-1}$. Then for any $n \in \mathbb{N}$, using (11) and $q_{n+1} = a_{n+1}q_n + q_{n-1} > a_{n+1}q_n$ we get

$$|q_n v(a) - p_n| < \frac{1}{q_{n+1}} < \frac{1}{a_{n+1}q_n} < \frac{1}{q_n^n},$$

hence $\|q_n v(a)\| < \frac{1}{q_n^n}$, and then

$$\sum_{j=1}^{q_n} \frac{1}{\|jv(a)\|} > \frac{1}{\|q_n v(a)\|} > q_n^n.$$

Using $\epsilon = 1$, it is then straightforward to check that there is no constant C such that $\sum_{j=1}^m \frac{1}{\|jv(a)\|} \leq Cm(\log m)^2$ for all m .

We will need the following lemma [19, p. 324, Lemma 3] to prove a theorem; cf. Khinchin [78, p. 63, Theorem 30].

$$m \mapsto \frac{1}{m \log m} \sum_{j=1}^m \frac{1}{\|jx\|} \text{ for } x = \frac{-1+\sqrt{5}}{2}$$




Figure 1: $\frac{1}{m \log m} \sum_{j=1}^m \frac{1}{\|jx\|}$ for $x = \frac{-1+\sqrt{5}}{2}$, for $m = 5000, \dots, 40000$

Lemma 12. *If ϕ is a function defined on the positive integers such that $\phi(n) \geq 1$ for all n and*

$$\sum_{n=1}^{\infty} \frac{1}{\phi(n)} < \infty,$$

then for almost all $x \in \Omega$ there are only finitely many n such that $a_n(x) \geq \phi(n)$.

Proof. For a measurable set $A \subset \Omega$, define

$$\gamma(A) = \frac{1}{\log 2} \int_A \frac{1}{1+x} d\mu(x),$$

in other words $d\gamma(x) = \frac{1}{(1+x)\log 2} d\mu(x)$. Thus for a measurable set $A \subset \Omega$ we have

$$\gamma(A) \leq \frac{1}{\log 2} \int_A dx = \frac{1}{\log 2} \mu(A)$$

and

$$\gamma(A) \geq \frac{1}{\log 2} \int_A \frac{1}{2} dx = \frac{1}{2\log 2} \mu(A).$$

We will use that γ is an invariant measure for the Gauss transformation $T : \Omega \rightarrow \Omega$ [44, p. 77, Lemma 3.5], i.e., if $A \subset \Omega$ is a measurable set then

$$(T_*\gamma)(A) = \gamma(T^{-1}(A)) = \gamma(A).$$

Let $A_n = \{x \in \Omega : a_n(x) \geq \phi(n)\}$, $n \geq 1$. As

$$a_n(x) = \left\lfloor \frac{1}{T^{n-1}(x)} \right\rfloor,$$

we have

$$\begin{aligned} A_n &\subset \left\{x \in \Omega : \frac{1}{T^{n-1}(x)} > \phi(n)\right\} \\ &= \left\{x \in \Omega : T^{n-1}(x) < \frac{1}{\phi(n)}\right\} \\ &= (T^{n-1})^{-1} \left(\left[0, \frac{1}{\phi(n)}\right) \setminus \mathbb{Q} \right). \end{aligned}$$

Hence

$$\begin{aligned} \mu(A_n) &\leq \mu \left((T^{n-1})^{-1} \left(\left[0, \frac{1}{\phi(n)}\right) \setminus \mathbb{Q} \right) \right) \\ &\leq 2\log 2 \cdot \gamma \left((T^{n-1})^{-1} \left(\left[0, \frac{1}{\phi(n)}\right) \setminus \mathbb{Q} \right) \right) \\ &= 2\log 2 \cdot \gamma \left(\left[0, \frac{1}{\phi(n)}\right) \setminus \mathbb{Q} \right) \\ &\leq 2\log 2 \cdot \frac{1}{\log 2} \cdot \mu \left(\left[0, \frac{1}{\phi(n)}\right) \setminus \mathbb{Q} \right) \\ &= \frac{2}{\phi(n)}. \end{aligned}$$

It follows that

$$\sum_{n=1}^{\infty} \mu(A_n) < \infty,$$

and thus by the Borel-Cantelli lemma [19, p. 59, Theorem 4.3] we have

$$\mu(\limsup_{n \rightarrow \infty} A_n) = 0.$$

□

Let λ be Lebesgue measure on $I = [0, 1]$, let $d\gamma(x) = \frac{1}{(1+x)\log 2} d\lambda(x)$, and let $T : I \rightarrow I$ be the Gauss transformation, $T(x) = x^{-1} - [x]^{-1}$ for $x > 0$ and $T(0) = 0$, for which $T_*\gamma = \gamma$ [44, p. 77, Lemma 3.5]. Suppose that ν is a Borel probability measure on $[0, 1]$ such that the pushforward measure $T_*\nu$ is absolutely continuous with respect to ν . For $f \in L^1(\nu)$, define $d\nu_f = f d\nu$, and define $P_\nu : L^1(\nu) \rightarrow L^1(\nu)$ by

$$P_\nu f = \frac{d(T_*\nu_f)}{d\nu}, \quad f \in L^1(\nu).$$

Thus, for $g \in L^\infty(\nu)$, using the change of variables formula,

$$\int_I g \cdot P_\nu f d\nu = \int_I g d(T_*\nu_f) = \int_I g \circ T d\nu_f = \int_I (g \circ T) \cdot f d\nu,$$

in particular,

$$\int_I P_\nu f d\nu = \int_I f d\nu.$$

We call $P_\nu : L^1(\nu) \rightarrow L^1(\nu)$ a **Perron-Frobenius operator for T** . It is a fact that if $f \geq 0$ then $P_\nu f \geq 0$ [69, p. 57, Proposition 2.1.1], namely $P_\nu \geq 0$. It can be proved that for $f \in L^1(\gamma)$, for almost all $x \in I$ [69, p. 59, Proposition 2.1.2],

$$(P_\gamma f)(x) = \sum_{k=1}^{\infty} \frac{x+1}{(x+k)(x+k+1)} \cdot f\left(\frac{1}{x+k}\right),$$

and for $f \in L^1(\lambda)$, for almost all $x \in I$ [69, p. 60, Corollary 2.1.4],

$$(P_\lambda f)(x) = \sum_{k=1}^{\infty} \frac{1}{(x+k)^2} \cdot f\left(\frac{1}{x+k}\right),$$

and with $g(x) = (x+1)f(x)$, for $n \geq 1$ it holds for almost all $x \in I$ that $(P_\lambda^n f)(x) = \frac{(P_\gamma^n g)(x)}{x+1}$. Iosifescu and Kraaikamp [69, Chapter 2] give a detailed presentation of Perron-Frobenius operators for the Gauss map. We make the final remark that $P_\nu 1_I = 1_I$ is equivalent with $\int_I 1_E d\nu = \int_I 1_{T^{-1}(E)} d\nu$ for all Borel sets E in I , i.e. $\nu(E) = \nu(T^{-1}(E))$, which in turn means $T_*\nu = \nu$, cf. Markov operators [83, Chapter 5, §5.1, pp. 177-186]. An object similar to Perron-Frobenius operators for the Gauss transformation is the **zeta-function for the Gauss transformation**, for which see Lagarias [87, p. 58, §3.3].

The following theorem gives a lower bound on the sum $\sum_{j=1}^m \frac{1}{\|jx\|}$, cf. [151, p. 4, Theorem 3.1].

Theorem 13. For almost all $x \in \Omega$ there is some $C > 0$ such that

$$\sum_{j=1}^m \frac{1}{\|jx\|} > Cm \log m.$$

Proof. For all $x \in \Omega$, if $n \geq 1$ then $q_n \geq 2^{\frac{n-1}{2}}$, by (9). Take $\phi(n) = 2^{\frac{n-2}{2}}$. The series $\sum_{n=1}^{\infty} \frac{1}{\phi(n)}$ converges, so by Lemma 12, for almost all $x \in \Omega$ there are only finitely many n such that $a_n \geq \phi(n)$. That is, for almost all $x \in \Omega$ there is some n_0 such that if $n \geq n_0$ then

$$a_n < \phi(n) = 2^{\frac{n-2}{2}} \leq q_{n-1}.$$

Hence, if $n \geq n_0$ then

$$q_n = a_n q_{n-1} + q_{n-2} < q_{n-1}^2 + q_{n-2} < 2q_{n-1}^2.$$

It follows that for almost all $x \in \Omega$ there is some K such that

$$q_{n+1} < Kq_n^2 \tag{15}$$

for all $n \geq 0$.

For such an x , let m be a positive integer and let $q_n \leq m < q_{n+1}$. For $1 \leq j \leq m$ we have by (11),

$$\left\| jx - \frac{jp_n}{q_n} \right\| \leq \left| jx - \frac{jp_n}{q_n} \right| = j \left| x - \frac{p_n}{q_n} \right| < \frac{j}{q_n q_{n+1}} < \frac{1}{q_n}.$$

Therefore for $1 \leq j \leq m$ we have

$$\|jx\| \leq \left\| jx - \frac{jp_n}{q_n} \right\| + \left\| \frac{jp_n}{q_n} \right\| < \frac{1}{q_n} + \left\| \frac{jp_n}{q_n} \right\|.$$

Let $L = \lfloor \frac{m}{q_n} \rfloor$, so $Lq_n \leq m$. Then,

$$\begin{aligned} \sum_{j=1}^m \frac{1}{\|jx\|} &> \sum_{j=1}^m \frac{1}{\frac{1}{q_n} + \left\| \frac{jp_n}{q_n} \right\|} \\ &\geq \sum_{l=0}^{L-1} \sum_{h=1}^{q_n} \frac{1}{\frac{1}{q_n} + \left\| \frac{(lq_n+h)p_n}{q_n} \right\|} \\ &= q_n \sum_{l=0}^{L-1} \sum_{h=1}^{q_n} \frac{1}{1 + q_n \left\| \frac{hp_n}{q_n} \right\|} \\ &= Lq_n \sum_{h=1}^{q_n} \frac{1}{1 + q_n \left\| \frac{hp_n}{q_n} \right\|} \\ &= Lq_n \sum_{k=0}^{q_n-1} \frac{1}{1 + q_n \cdot \frac{k}{q_n}} \\ &> Lq_n \log q_n. \end{aligned}$$

But if $y \geq 1$ then $[y] > \frac{y}{2}$, so $L = [\frac{m}{q_n}] > \frac{m}{2q_n}$. Hence by (15),

$$\sum_{j=1}^m \frac{1}{\|jx\|} > \frac{m}{2} \log q_n > \frac{m}{2} \log \sqrt{\frac{q_{n+1}}{K}} > \frac{m}{2} \log \sqrt{\frac{m}{K}},$$

and thus there is some $C > 0$ such that $\sum_{j=1}^m \frac{1}{\|jx\|} > Cm \log m$ for all $m \geq 1$. \square

The following is from Kuipers and Niederreiter [85, p. 131, Exercise 3.12].

Theorem 14. *If $x \in \Omega$ is of type $< \psi$, then for all $m \geq 1$ we have*

$$\sum_{j=1}^m \frac{1}{j \|jx\|} < 24 \left((\log m)^2 + \psi(m) + \sum_{j=1}^m \frac{\psi(j)}{j} \right).$$

Proof. **Summation by parts** is the following identity, which can be easily checked:

$$\sum_{n=1}^N a_n (b_{n+1} - b_n) = a_{N+1} b_{N+1} - a_1 b_1 - \sum_{n=1}^N b_{n+1} (a_{n+1} - a_n).$$

Let $a_j = \frac{1}{j}$, let $s_j = \sum_{h=1}^j \frac{1}{\|hx\|}$, let $b_1 = 0$, and let $b_j = s_{j-1}$ for $j \geq 2$. Doing summation by parts gives

$$\sum_{j=1}^m \frac{1}{j \|jx\|} = \frac{1}{m+1} s_m - \sum_{j=1}^m s_j \left(\frac{1}{j+1} - \frac{1}{j} \right) = \frac{1}{m+1} s_m + \sum_{j=1}^m s_j \frac{1}{j(j+1)}.$$

As x is of type $< \psi$, we can use Theorem 10 to get $s_j < 12j(\psi(j) + \log j)$ for each $j \geq 1$. Therefore

$$\begin{aligned} \sum_{j=1}^m \frac{1}{j \|jx\|} &< \frac{1}{m+1} 12m(\psi(m) + \log m) + \sum_{j=1}^m \frac{12(\psi(j) + \log j)}{j+1} \\ &< 12(\psi(m) + \log m) + 12(\log m)^2 + 12 \sum_{j=1}^m \frac{\psi(j)}{j+1} \\ &< 24(\log m)^2 + 12\psi(m) + 12 \sum_{j=1}^m \frac{\psi(j)}{j+1}. \end{aligned}$$

\square

Erdős [45] proves that for almost all x ,

$$\sum_{j=1}^m \frac{1}{j \|jx\|} = (1 + o(1))(\log m)^2.$$

Kruse [84] gives a comprehensive investigation of the sums $\sum_{j=1}^m \frac{1}{j^s \|jx\|^t}$, $s, t \geq 0$. The results depend on whether s and t are less than, equal, or greater than 1, and on whether $t < s$. One of the theorems proved by Kruse is the following [84, p. 260, Theorem 7]. If $t > 1$ and $0 \leq s \leq t$, and if $\epsilon > 0$, then for almost all x we have

$$\sum_{j=1}^m \frac{1}{j^s \|jx\|^t} = O\left(m^{t-s} (\log m)^{(1+\epsilon)t}\right).$$

Haber and Osgood [56, p. 387, Theorem 1] prove that for real $t \geq 1$, $A > 1$, $M > 0$, $r > 0$, there is some $C = C(t, A, M, r) > 0$ such that for all $x \in \Omega$ satisfying $q_{n+1}(x) < Mq_n(x)^r$, for all positive integers K ,

$$\sum_{n=K+1}^{[AK]} \|nx\|^{-t} > \begin{cases} CK \log K & t = 1 \\ CK^{1+(t-1)/r} & t > 1. \end{cases}$$

We remind ourselves that according to Theorem 4, the elements of $\mathcal{D}(r+1)$ are those $x \in \Omega$ for which there is some $c(x) > 0$ such that $q_{n+1}(x) \leq C(x)q_n(x)^r$ for all $n \geq 1$.

For $x \in \mathbb{Z} + \frac{1}{2}$, define $\{\{x\}\} = \frac{1}{2}$. If $x \notin \mathbb{Z} + \frac{1}{2}$, then there is an integer m_x for which $|x - m_x| < |x - n|$ for all integers $n \neq m_x$, and we define $\{\{x\}\} = x - m_x$. Sinai and Ulcigrai [139, p. 96, Proposition 2] prove that if α has bounded partial quotients, then there is some $C(\alpha)$ such that for all M ,

$$\left| \sum_{m=1}^M \frac{1}{\{\{m\alpha\}\}} \right| \leq C(\alpha)M.$$

7 Weyl's inequality, Vinogradov's estimate, Farey fractions, and the circle method

Write

$$\mathcal{A} = \{(a, q) \in \mathbb{Z}^2 : \gcd(a, q) = 1, q \geq 1\}.$$

We first prove four estimates following Nathanson [107, pp. 104–110, Lemmas 4.8–4.11] that we will use in what follows; cf. Vinogradov [150, p. 26, Chapter I, Lemma 8b].

Lemma 15. *There is some C such that if $\alpha \in \mathbb{R}$, $(a, q) \in \mathcal{A}$, and*

$$\left| \alpha - \frac{a}{q} \right| \leq \frac{1}{q^2},$$

then

$$\sum_{1 \leq r \leq q/2} \frac{1}{\|\alpha r\|} \leq Cq \log q.$$

Proof. For $q = 1$, $\sum_{1 \leq r \leq q/2} \frac{1}{\|ar\|} = 0$. For $q \geq 2$, let $1 \leq r \leq \frac{q}{2}$. As $\gcd(a, q) = 1$ and $r \not\equiv 0 \pmod{q}$, $ar \not\equiv 0 \pmod{q}$. So for $\mu_r = \left\lfloor \frac{ar}{q} \right\rfloor$, there is some $1 \leq \sigma_r \leq q - 1$ such that $ar = \mu_r q + \sigma_r$. Then

$$\left\| \frac{ar}{q} \right\| = \left\| \frac{\sigma_r}{q} \right\| \in \left\{ \frac{\sigma_r}{q}, 1 - \frac{\sigma_r}{q} \right\} = \left\{ \frac{\sigma_r}{q}, \frac{q - \sigma_r}{q} \right\}.$$

Put $\frac{s_r}{q} = \left\| \frac{ar}{q} \right\|$, so (i) $s_r = \sigma_r$ or (ii) $s_r = q - \sigma_r$. In case (i), $\frac{s_r}{q} = \frac{ar}{q} - \mu_r$. In case (ii), $\frac{s_r}{q} = 1 - \left(\frac{ar}{q} - \mu_r \right)$. In case (i) let $\epsilon_r = 1, m_r = \mu_r$, and in case (ii) let $\epsilon_r = -1, m_r = \mu_r + 1$. Thus, whether (i) or (ii) holds we have

$$\frac{s_r}{q} = \epsilon_r \left(\frac{ar}{q} - m_r \right), \quad \frac{s_r}{q} = \left\| \frac{ar}{q} \right\|, \quad 1 \leq s_r \leq \frac{q}{2}.$$

Write

$$\alpha - \frac{a}{q} = \frac{\theta}{q^2},$$

for some real θ , $|\theta| \leq 1$. For $\theta_r = \frac{2r}{q}\theta$, which satisfies $|\theta_r| \leq |\theta| \leq 1$,

$$\alpha r = \frac{ar}{q} + \frac{r\theta}{q^2} = \frac{ar}{q} + \frac{\theta_r}{2q}.$$

Then

$$\begin{aligned} \|\alpha r\| &= \left\| \frac{ar}{q} + \frac{\theta_r}{2q} \right\| \\ &= \left\| \epsilon_r \frac{s_r}{q} + m_r + \frac{\theta_r}{2q} \right\| \\ &\geq \left\| \epsilon_r \frac{s_r}{q} + m_r \right\| - \left\| \frac{\theta_r}{2q} \right\| \\ &= \frac{s_r}{q} - \left| \frac{\theta_r}{2q} \right| \\ &\geq \frac{s_r}{q} - \frac{1}{2q}. \end{aligned}$$

Take $1 \leq r_1, r_2 \leq \frac{q}{2}$ and suppose that $s_{r_1} = s_{r_2}$. So

$$\epsilon_{r_1} \left(\frac{ar_1}{q} - m_{r_1} \right) = \epsilon_{r_2} \left(\frac{ar_2}{q} - m_{r_2} \right)$$

hence $ar_1 \equiv \epsilon_{r_1} \epsilon_{r_2} ar_2 \pmod{q}$. As $\gcd(a, q) = 1$, $r_1 \equiv \epsilon_{r_1} \epsilon_{r_2} r_2 \pmod{q}$. Because $1 \leq r_1, r_2 \leq q$, if $r_1 \equiv r_2 \pmod{q}$ then $r_1 = r_2$ and if $r_1 \equiv -r_2 \pmod{q}$ then $r_1 = \frac{q}{2}$ and $r_2 = \frac{q}{2}$, so in any case $r_1 = r_2$. Therefore

$$\left\{ \frac{s_r}{q} : 1 \leq r \leq \frac{q}{2} \right\} = \left\{ \frac{s}{q} : 1 \leq s \leq \frac{q}{2} \right\}.$$

Using the two things we have established,

$$\begin{aligned}
\sum_{1 \leq r \leq q/2} \frac{1}{\|\alpha r\|} &\leq \sum_{1 \leq r \leq q/2} \frac{1}{\frac{s_r}{q} - \frac{1}{2q}} \\
&= \sum_{1 \leq s \leq q/2} \frac{1}{\frac{s}{q} - \frac{1}{2q}} \\
&= 2q \sum_{1 \leq s \leq q/2} \frac{1}{2s-1} \\
&\leq 2q \sum_{1 \leq s \leq q/2} \frac{1}{s} \\
&\leq 2q \left(\log \frac{q}{2} + \gamma + O(q^{-1}) \right) \\
&= O(q \log q).
\end{aligned}$$

□

Lemma 16. *There is some C such that if $\alpha \in \mathbb{R}$, $(a, q) \in \mathcal{A}$, and*

$$\left| \alpha - \frac{a}{q} \right| \leq \frac{1}{q^2},$$

then for any positive real V and nonnegative integer h ,

$$\sum_{r=1}^q \min \left(V, \frac{1}{\|\alpha(hq+r)\|} \right) \leq C(V + q \log q).$$

Proof. Write

$$\alpha = \frac{a}{q} + \frac{\theta}{q^2},$$

which satisfies $|\theta| \leq 1$, and for $1 \leq r \leq q$ define

$$\delta_r = R(\theta h) + \frac{\theta r}{q},$$

which satisfies $-1 \leq \delta_r < 2$. Then

$$\begin{aligned}
\alpha(hq+r) &= \left(\frac{a}{q} + \frac{\theta}{q^2} \right) (hq+r) \\
&= ah + \frac{ar}{q} + \frac{\theta h}{q} + \frac{\theta r}{q^2} \\
&= ah + \frac{ar}{q} + \frac{R(\theta h) + [\theta h]}{q} + \frac{\theta r}{q^2} \\
&= ah + \frac{ar + [\theta h] + \delta_r}{q}.
\end{aligned}$$

For $m_r = \left\lceil \frac{ar + [\theta h] + \delta_r}{q^2} \right\rceil$,

$$R(\alpha(hq + r)) = R\left(\frac{ar + [\theta h] + \delta_r}{q}\right) = \frac{ar + [\theta h] + \delta_r}{q} - m_r.$$

Suppose that $t \in \left[0, 1 - \frac{1}{q}\right]$ and that $t \leq R(\alpha(hq + r)) \leq t + \frac{1}{q}$. Then

$$qt \leq ar + [\theta h] + \delta_r - qm_r \leq qt + 1.$$

This implies, as $\delta_r \geq -1$,

$$ar - qm_r \leq qt + 1 - [\theta h] - \delta_r \leq qt + 1 - [\theta h] + 1 = qt - [\theta h] + 2$$

and, as $\delta_r < 2$,

$$ar - qm_r \geq qt - [\theta h] - \delta_r > qt - [\theta h] - 2,$$

so $ar - qm_r \in J_t$, writing

$$J_t = (qt - [\theta h] - 2, qt - [\theta h] + 2).$$

For $1 \leq r_1, r_2 \leq q$, if $ar_1 - qm_{r_1} = ar_2 - qm_{r_2}$ then $ar_1 \equiv ar_2 \pmod{q}$, and $\gcd(a, q) = 1$ implies $r_1 \equiv r_2 \pmod{q}$; and $1 \leq r_1, r_2 \leq q$ so $r_1 = r_2$. For $t \in \left[0, 1 - \frac{1}{q}\right]$, four integers belong to J_t , hence

$$\{1 \leq r \leq q : ar - qm_r \in J_t\}$$

has at most four elements. But

$$\left\{1 \leq r \leq q : R(\alpha(hq + r)) \in \left[t, t + \frac{1}{q}\right]\right\} \subset \{1 \leq r \leq q : ar - qm_r \in J_t\}.$$

Now,

$$\begin{aligned} & \left\{1 \leq r \leq q : \|\alpha(hq + r)\| \in \left[t, t + \frac{1}{q}\right]\right\} \\ &= \left\{1 \leq r \leq q : R(\alpha(hq + r)) \in \left[t, t + \frac{1}{q}\right]\right\} \\ & \cup \left\{1 \leq r \leq q : 1 - R(\alpha(hq + r)) \in \left[t, t + \frac{1}{q}\right]\right\} \\ &= \left\{1 \leq r \leq q : R(\alpha(hq + r)) \in \left[t, t + \frac{1}{q}\right]\right\} \\ & \cup \left\{1 \leq r \leq q : R(\alpha(hq + r)) \in \left[1 - \frac{1}{q} - t, 1 - t\right]\right\}, \end{aligned}$$

whence

$$\begin{aligned} \left\{1 \leq r \leq q : \|\alpha(hq + r)\| \in \left[t, t + \frac{1}{q}\right]\right\} & \subset \{1 \leq r \leq q : ar - qm_r \in J_t\} \\ & \cup \{1 \leq r \leq q : ar - qm_r \in J_{1 - \frac{1}{q} - t}\}. \end{aligned}$$

This shows that if $t \in \left[0, 1 - \frac{1}{q}\right]$ then

$$\left\{1 \leq r \leq q : \|\alpha(hq + r)\| \in \left[t, t + \frac{1}{q}\right]\right\}$$

has at most eight elements. For $0 \leq k < \frac{q}{2}$, writing

$$I_k = \left[\frac{k}{q}, \frac{k}{q} + \frac{1}{q}\right],$$

the set $\{1 \leq r \leq q : \|\alpha(hq + r)\| \in I_k\}$ has at most eight elements. Therefore

$$\begin{aligned} \sum_{1 \leq r \leq q} \min\left(V, \frac{1}{\|\alpha(hq + r)\|}\right) &= \sum_{0 \leq k < q/2} \sum_{1 \leq r \leq q, \|\alpha(hq + r)\| \in I_k} \min\left(V, \frac{1}{\|\alpha(hq + r)\|}\right) \\ &\leq 8V + \sum_{1 \leq k < q/2} \sum_{1 \leq r \leq q, \|\alpha(hq + r)\| \in I_k} \frac{1}{\|\alpha(hq + r)\|} \\ &\leq 8V + \sum_{1 \leq k < q/2} 8 \cdot \frac{q}{k} \\ &= O(V + q \log q). \end{aligned}$$

□

Lemma 17. *There is some C such that if $\alpha \in \mathbb{R}$, $(a, q) \in \mathcal{A}$,*

$$\left|\alpha - \frac{a}{q}\right| \leq \frac{1}{q^2},$$

$U \geq 1$ is a real number, and n is a positive integer, then

$$\sum_{1 \leq k \leq U} \min\left(\frac{n}{k}, \frac{1}{\|\alpha k\|}\right) \leq C \left(\frac{n}{q} + U + q\right) \log 2qU.$$

Proof. For $1 \leq k \leq U$ there is some $0 \leq h_k < \frac{U}{q}$ and $1 \leq r_k \leq q$ such that $k = qh_k + r_k$, and then

$$\begin{aligned} \sum_{1 \leq k \leq U} \min\left(\frac{n}{k}, \frac{1}{\|\alpha k\|}\right) &\leq \sum_{0 \leq h < U/q} \sum_{1 \leq r \leq q} \min\left(\frac{n}{qh + r}, \frac{1}{\|\alpha(hq + r)\|}\right) \\ &\leq \sum_{1 \leq r \leq q/2} \frac{1}{\|\alpha r\|} + \sum_{q/2 < r \leq q} \min\left(\frac{n}{r}, \frac{1}{\|\alpha r\|}\right) \\ &\quad + \sum_{1 \leq h < U/q} \sum_{1 \leq r \leq q} \min\left(\frac{n}{qh + r}, \frac{1}{\|\alpha(hq + r)\|}\right) \\ &\leq C_1 q \log q + \sum_{q/2 < r \leq q} \min\left(\frac{n}{r}, \frac{1}{\|\alpha r\|}\right) \\ &\quad + \sum_{1 \leq h < U/q} \sum_{1 \leq r \leq q} \min\left(\frac{n}{qh + r}, \frac{1}{\|\alpha(hq + r)\|}\right), \end{aligned}$$

the last inequality by Lemma 15. If $\frac{q}{2} < r \leq q$ then $\frac{1}{r} < \frac{2}{q} = \frac{2}{(h+1)q}$ for $h = 0$, and if $1 \leq h < \frac{U}{q}$ and $1 \leq r \leq q$ then $h \geq \frac{h+1}{2}$ so $hq + r > hq \geq \frac{(h+1)q}{2}$ and hence $\frac{1}{hq+r} < \frac{2}{(h+1)q}$, whence

$$\begin{aligned} & \sum_{q/2 < r \leq q} \min\left(\frac{n}{r}, \frac{1}{\|\alpha r\|}\right) + \sum_{1 \leq h < U/q} \sum_{1 \leq r \leq q} \min\left(\frac{n}{qh+r}, \frac{1}{\|\alpha(hq+r)\|}\right) \\ \leq & 2 \sum_{q/2 < r \leq q} \min\left(\frac{n}{(h+1)q}, \frac{1}{\|\alpha r\|}\right) + 2 \sum_{1 \leq h < U/q} \sum_{1 \leq r \leq q} \min\left(\frac{n}{(h+1)q}, \frac{1}{\|\alpha(hq+r)\|}\right). \end{aligned}$$

Consequently

$$\sum_{1 \leq k \leq U} \min\left(\frac{n}{k}, \frac{1}{\|\alpha k\|}\right) \leq C_1 q \log q + 2 \sum_{0 \leq h < U/q} \sum_{1 \leq r \leq q} \min\left(\frac{n}{(h+1)q}, \frac{1}{\|\alpha(hq+r)\|}\right).$$

Lemma 16 with $V = \frac{n}{(h+1)q}$ says

$$\sum_{1 \leq r \leq q} \min\left(\frac{n}{(h+1)q}, \frac{1}{\|\alpha(hq+r)\|}\right) \leq C_2 \left(\frac{n}{(h+1)q} + q \log q\right),$$

therefore

$$\begin{aligned} \sum_{1 \leq k \leq U} \min\left(\frac{n}{k}, \frac{1}{\|\alpha k\|}\right) & \ll q \log q + \sum_{0 \leq h < U/q} \left(\frac{n}{(h+1)q} + q \log q\right) \\ & \ll q \log q + \frac{n}{q} \sum_{1 \leq h < \frac{U}{q} + 1} \frac{1}{h} + q(\log q) \left(\frac{U}{q} + 1\right) \\ & \ll q \log q + \frac{n}{q} \log\left(\frac{U}{q} + 1\right) + U \log q \\ & \ll U \log 2qU + q \log 2qU + \frac{n}{q} \log\left(\frac{U}{q} + 1\right). \end{aligned}$$

If $U \leq q$ then $\frac{U}{q} + 1 \leq 2 \leq 2qU$, and if $U > q$ then $\frac{U}{q} + 1 \leq U + 1 \leq 2U \leq 2qU$, hence

$$\sum_{1 \leq k \leq U} \min\left(\frac{n}{k}, \frac{1}{\|\alpha k\|}\right) \ll U \log 2qU + q \log 2qU + \frac{n}{q} \log 2qU.$$

□

Lemma 18. *There is some C such that if $\alpha \in \mathbb{R}$, $(a, q) \in \mathcal{A}$,*

$$\left|\alpha - \frac{p}{q}\right| \leq \frac{1}{q^2},$$

and $U, V \geq 1$ are real numbers, then

$$\sum_{1 \leq k \leq U} \min \left(V, \frac{1}{\|\alpha k\|} \right) \leq C \left(q + U + V + \frac{UV}{q} \right) \max\{1, \log q\}.$$

Proof. For $1 \leq k \leq U$ there is some $0 \leq h_k < \frac{U}{q}$ and $1 \leq r_k \leq q$ such that $k = qh_k + r_k$, and then, as in the proof of Lemma 17,

$$\begin{aligned} \sum_{1 \leq k \leq U} \min \left(V, \frac{1}{\|\alpha k\|} \right) &\leq \sum_{0 \leq h < U/q} \sum_{1 \leq r \leq q} \min \left(V, \frac{1}{\|\alpha(hq + r)\|} \right) \\ &\leq C_1 q \log q + 2 \sum_{0 \leq h < U/q} \sum_{1 \leq r \leq q} \min \left(V, \frac{1}{\|\alpha(hq + r)\|} \right). \end{aligned}$$

Using Lemma 16,

$$\begin{aligned} \sum_{1 \leq k \leq U} \min \left(V, \frac{1}{\|\alpha k\|} \right) &\leq C_1 q \log q + 2C_2 \sum_{0 \leq h < U/q} (V + q \log q) \\ &\ll q \log q + (V + q \log q) \left(\frac{U}{q} + 1 \right) \\ &\ll q \log q + \frac{UV}{q} + V + U \log q. \end{aligned}$$

□

Weyl's inequality [107, p. 114, Theorem 4.3] is the following. For $k \geq 2$ and $\epsilon > 0$, there is some $C(k, \epsilon)$ such that if $\alpha \in \mathbb{R}$, $f(x)$ is a real polynomial with highest degree term αx^k , $(a, q) \in \mathcal{A}$, and

$$\left| \alpha - \frac{a}{q} \right| \leq \frac{1}{q^2},$$

then, writing $S_N(f) = \sum_{j=1}^N e^{2\pi i f(j)}$ and $K = 2^{k-1}$,

$$|S_N(f)| \leq C(k, \epsilon) \cdot N^{1+\epsilon} (N^{-1} + q^{-1} + N^{-k} q)^{\frac{1}{K}}.$$

Weyl's inequality is proved using Lemma 18.

Montgomery [102, Chapter 3] gives a similar but more streamlined presentation of Weyl's inequality. Chandrasekharan [29] gives a historical survey of exponential sums.

Vinogradov's estimate [147, p. 26, Theorem 3.1] states that there is some C such that for $n \geq 2$, $1 \leq q \leq n$, $\gcd(a, q) = 1$, and $\left| \alpha - \frac{a}{q} \right| \leq q^{-2}$, then

$$|f_n(\alpha)| \leq C(nq^{-1/2} + n^{4/5} + n^{1/2}q^{1/2})(\log n)^4, \quad (16)$$

where $f_n(\alpha) = \sum_{p \leq n} (\log p) e^{2\pi i \alpha p}$; cf. Nathanson [107, p. 220, Theorem 8.5] and Vinogradov [150, p. 131, Chapter IX, Theorem 1]. This is proved using Lemma 17.

Fix $B > 0$ and let $P_n = (\log n)^B$. For $1 \leq a \leq q \leq P_n$ and $\gcd(a, q) = 1$, let

$$\mathfrak{M}_n(q, a) = \left\{ \alpha \in \mathbb{R} : \left| \alpha - \frac{a}{q} \right| \leq P_n n^{-1} \right\},$$

called a **major arc**. One checks that there is some n_B such that if $n \geq n_B$, then $\mathfrak{M}_n(q, a)$ and $\mathfrak{M}_n(q', a')$ are disjoint when $(q, a) \neq (q', a')$. Let

$$\mathfrak{M}_n = \bigcup_{1 \leq a \leq q \leq P_n, \gcd(a, q) = 1} \mathfrak{M}_n(q, a).$$

The **Farey fractions of order N** are

$$\mathfrak{F}_N = \left\{ \frac{h}{k} : 0 \leq h \leq k \leq N, \gcd(h, k) = 1 \right\}.$$

Cf. the **Stern-Brocot tree** [55, §4.5]. For early appearances of Farey fractions, see Dickson [35, pp. 155–158, Chapter V]. It is proved by Cauchy that if h/k and h'/k' are successive elements of \mathfrak{F}_N , then $kh' - hk' = 1$ [61, p. 23, Theorem 28]. Let

$$\phi(m) = |\{1 \leq k \leq m : \gcd(k, m) = 1\}|,$$

the **Euler phi function**, and write $\Phi(N) = \sum_{1 \leq m \leq N} \phi(m)$. One sees that $|\mathfrak{F}_N| = 1 + \Phi(N)$, and it was proved by Mertens [61, p. 268] that

$$\Phi(N) = \frac{3N^2}{\pi^2} + O(N \log N).$$

Let λ be Lebesgue measure on \mathbb{R} . For $n \geq n_B$, because the major arcs are pairwise disjoint,

$$\begin{aligned} \lambda(\mathfrak{M}_n) &= \sum_{1 \leq a \leq q \leq P_n, \gcd(a, q) = 1} 2P_n n^{-1} \\ &= \left(\sum_{m=1}^{P_n} \phi(m) \right) 2P_n n^{-1} \\ &= \frac{6}{\pi^2} P_n^3 n^{-1} + O(P_n^2 n^{-1} \log P_n). \end{aligned}$$

Let $I_n = (P_n n^{-1}, 1 + P_n n^{-1}]$, which for $n > 2P_n^2$ contains \mathfrak{M}_n . Let

$$\mathfrak{m}_n = I_n \setminus \mathfrak{M}_n,$$

called the **minor arcs**.

With $f_n(\alpha) = \sum_{p \leq n} (\log p) e^{2\pi i \alpha p}$ and

$$R(n) = \sum_{p_1 + p_2 + p_3 = n} (\log p_1)(\log p_2)(\log p_3),$$

we have

$$R(n) = \int_{I_n} f_n(\alpha)^3 e^{-2\pi i n \alpha} d\alpha = \int_{\mathfrak{M}_n} f_n(\alpha)^3 e^{-2\pi i n \alpha} d\alpha + \int_{\mathfrak{m}_n} f_n(\alpha)^3 e^{-2\pi i n \alpha} d\alpha.$$

Using (16), it can be proved that for $A > 0$ with $B \geq 2A + 10$ [147, p. 29, Theorem 3.2],

$$\int_{\mathfrak{m}_n} |f_n(\alpha)|^3 d\alpha = O(n^2 (\log n)^{-A}).$$

Writing

$$\mathfrak{S}(n) = \left(\prod_{p \nmid n} (1 + (p-1)^{-3}) \right) \prod_{p|n} (1 - (p-1)^{-2}),$$

called the **singular series**, it is proved, using the Siegel-Walfisz theorem on primes in arithmetic progressions [103, p. 381, Corollary 11.19], that for $A > 0$ with $B \geq 2A$ [147, p. 31, Theorem 3.3],

$$\int_{\mathfrak{M}_n} f_n(\alpha)^3 e^{-2\pi i n \alpha} d\alpha = \frac{1}{2} n^2 \mathfrak{S}(n) + O(n^2 (\log n)^{-A}).$$

Thus

$$R(n) = \frac{1}{2} n^2 \mathfrak{S}(n) + O(n^2 (\log n)^{-A}),$$

and it follows from this that there is some n_0 such that if $n \geq n_0$ is odd then there are primes p_1, p_2, p_3 such that $n = p_1 + p_2 + p_3$.

For integers a, b with $\gcd(a, b) = 1$, the **Ford circle** $C(a, b)$ is the circle in \mathbb{C} that touches the line $\text{Im } z = 0$ at $z = \frac{a}{b}$ and has radius $\frac{1}{2b^2}$; in other words, $C(a, b)$ is the circle in \mathbb{C} with center $\frac{a}{b} + \frac{i}{2b^2}$ and radius $\frac{1}{2b^2}$. It is straightforward to prove that if $C(a, b)$ and $C(c, d)$ are Ford circles, then they are tangent if and only if $(bc - ad)^2 = 1$, and otherwise they are disjoint [4, p. 100, Theorem 5.6]. It is also straightforward to prove [4, p. 101, Theorem 5.7] that if $\frac{h_1}{k_1} < \frac{h}{k} < \frac{h_2}{k_2}$ are successive elements of \mathfrak{F}_N , then $C(h_1, k_1)$ and $C(h, k)$ touch at

$$\frac{h}{k} - \frac{k_1}{k(k^2 + k_1^2)} + \frac{i}{k^2 + k_1^2}$$

and $C(h, k)$ and $C(h_2, k_2)$ touch at

$$\frac{h}{k} + \frac{k_2}{k(k^2 + k_2^2)} + \frac{i}{k^2 + k_2^2}.$$

Bonahon [21, pp. 207 ff., Chapter 8] explains Ford circles in the language of hyperbolic geometry.

We remind ourselves that $|\mathfrak{F}_N| = 1 + \Phi(N) = 1 + \sum_{m=1}^N \phi(m)$ and let $\rho_{0,N} < \dots < \rho_{\Phi(N),N}$ be the elements of \mathfrak{F}_N . In particular, $\rho_{0,N} = 0$ and $\rho_{\Phi(N),N} = 1$. For $\rho_{n,N} = \frac{h_{n,N}}{k_{n,N}}$ with $\gcd(h_{n,N}, k_{n,N}) = 1$, write $C_{n,N} = C(h_{n,N}, k_{n,N})$.

Let $A_{0,N}$ be the clockwise arc of $C_{0,N}$ from i to the point at which $C_{0,N}$ and $C_{1,N}$ touch. For $0 < n < \Phi(N)$, let $A_{n,N}$ be the clockwise arc of $C_{n,N}$ from the point at which $C_{n-1,N}$ and $C_{n,N}$ touch to the point at which $C_{n,N}$ and $C_{n+1,N}$ touch. Finally, let $A_{\Phi(N),N}$ be the clockwise arc of $C_{\Phi(N),N}$ from the point at which $C_{\Phi(N)-1,N}$ and $C_{\Phi(N),N}$ touch to $i + 1$. Let A_N be the composition of the arcs

$$A_{0,N}, \dots, A_{\Phi(N),N}, \quad (17)$$

which is a contour from i to $i + 1$.

Write $H = \{\tau \in \mathbb{C} : \text{Im } \tau > 0\}$. The **Dedekind eta function** $\eta : H \rightarrow \mathbb{C}$ is defined by

$$\eta(\tau) = e^{\pi i \tau / 12} \prod_{m=1}^{\infty} (1 - e^{2\pi i m \tau}).$$

It is straightforward to check that η is analytic and that $\eta(\tau) \neq 0$ for all $\tau \in H$ [144, pp. 17–18, §1.44]. For $(h, k) \in \mathcal{A}$, let

$$s(h, k) = \sum_{r=1}^{k-1} \frac{r}{k} \left(\frac{hr}{k} - \left[\frac{hr}{k} \right] - \frac{1}{2} \right) = \sum_{r=1}^{k-1} \frac{r}{k} P_1(hr/k),$$

called a **Dedekind sum**; P_1 is the periodic Bernoulli function. Also, for $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbb{Z})$ write

$$\epsilon(a, b, c, d) = \exp \left(\pi i \left(\frac{a+d}{12c} + s(-d, c) \right) \right).$$

The **functional equation for the Dedekind eta function** [4, p. 52, Theorem 3.4] is

$$\eta \left(\frac{a\tau + b}{c\tau + d} \right) = \epsilon(a, b, c, d) (-i(c\tau + d))^{1/2} \eta(\tau), \quad \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbb{Z}), \tau \in H.$$

Let $p(n)$ be the number of ways of writing n as a sum of positive integers where the order does not matter, called the **partition function**. For example, $4, 3+1, 2+2, 2+1+1, 1+1+1+1$ are the partitions of 4, so $p(4) = 5$. Denoting by $D(0, 1)$ the open disc with center 0 and radius 1, define $F : D(0, 1) \rightarrow \mathbb{C}$ by

$$F(z) = \prod_{m=1}^{\infty} (1 - z^m)^{-1} = \sum_{n=0}^{\infty} p(n) z^n;$$

that the product and the series are equal was found by Euler. F is analytic.

On the one hand, $p(n) = \frac{F^{(n)}(0)}{n!}$, and on the other hand, by Cauchy's integral formula [144, p. 82, Theorem 2.41], if C is a circle with center 0 and radius $0 < R < 1$ then

$$F^{(n)}(0) = \frac{n!}{2\pi i} \int_C \frac{F(z)}{z^{n+1}} dz.$$

Taking C to be the circle with center 0 and radius $e^{-2\pi}$ and doing the change of variable $z = e^{2\pi i\tau}$,

$$\begin{aligned} p(n) &= \frac{1}{2\pi i} \int_i^{i+1} \frac{F(e^{2\pi i\tau})}{e^{2\pi i(n+1)\tau}} \cdot 2\pi i e^{2\pi i\tau} d\tau \\ &= \int_i^{i+1} F(e^{2\pi i\tau}) e^{-2\pi in\tau} d\tau \\ &= \int_{A_N} F(e^{2\pi i\tau}) e^{-2\pi in\tau} d\tau, \end{aligned}$$

where we remind ourselves that A_N is the contour (17) from i to $i+1$. Using this and the functional equation for the Dedekind eta function, Rademacher [4, p. 104, Theorem 5.10] proves that for $n \geq 1$,

$$p(n) = \frac{1}{\pi\sqrt{2}} \sum_{k=1}^{\infty} A_k(n) k^{1/2} \frac{d}{dn} \frac{\sinh\left(\pi\sqrt{\frac{2}{3}} \cdot \frac{1}{k} \sqrt{n - \frac{1}{24}}\right)}{\sqrt{n - \frac{1}{24}}},$$

for $A_k(n) = \sum_{0 \leq h < k, \gcd(h,k)=1} e^{\pi i s(h,k) - 2\pi i n h/k}$.

We make a final remark about the Farey fractions. Writing the elements of \mathfrak{F}_N as $\rho_{0,N} < \dots < \rho_{\Phi(N),N}$, let $\eta_{n,N} = \rho_{n,N} - \frac{n}{\Phi(N)}$ for $1 \leq n \leq N$. For example, $\Phi(5) = 10$ and

$$\{\rho_{1,5}, \dots, \rho_{10,5}\} = \left\{ \frac{1}{5}, \frac{1}{4}, \frac{1}{3}, \frac{2}{5}, \frac{1}{2}, \frac{3}{5}, \frac{2}{3}, \frac{3}{4}, \frac{4}{5}, 1 \right\},$$

and

$$\{\eta_{1,5}, \dots, \eta_{10,5}\} = \left\{ \frac{1}{10}, \frac{1}{20}, \frac{1}{30}, 0, 0, 0, -\frac{1}{30}, -\frac{1}{20}, -\frac{1}{10}, 0 \right\}.$$

Landau [91], following work of Franel, proves that the **Riemann hypothesis** is true if and only if for every $\epsilon > 0$,

$$\sum_{n=1}^{\Phi(N)} |\eta_{n,N}| = O(N^{\frac{1}{2} + \epsilon}).$$

For example, for $N = 5$, the left-hand side is $\frac{11}{30}$. See Narkiewicz [106, p. 40, §2.2.3].

8 Discrepancy and exponential sums

Discrepancy and Diophantine approximation are covered by Kuipers and Niederreiter [85, Chapters 1–2] and by Drmota and Tichy [41, §§1.1–1.4], especially [41, pp. 48–66, §1.4.1].

Let $\omega = (x_n)$, $n \geq 1$, be a sequence of real numbers, and let $E \subset [0, 1)$. For a positive integer N , let $A(E; N; \omega)$ be the number of x_n , $1 \leq n \leq N$, such that

$R(x_n) \in E$. We say that the sequence ω is **uniformly distributed modulo 1** if we have for all a and b with $0 \leq a < b \leq 1$ that

$$\lim_{N \rightarrow \infty} \frac{A([a, b]; N; \omega)}{N} = b - a.$$

It can be shown [85, p. 3, Corollary 1.1] that a sequence x_n is uniformly distributed modulo 1 if and only if for every Riemann integrable function $f : [0, 1] \rightarrow \mathbb{R}$ we have

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N f(R(x_n)) = \int_0^1 f(t) dt. \quad (18)$$

Thus, if a sequence is uniformly distributed then the integral of any Riemann integrable function on $[0, 1]$ can be approximated by sampling according to this sequence. This approximation can be quantified using the notion of discrepancy.

It can be proved [85, p. 7, Theorem 2.1] that a sequence x_n is uniformly distributed modulo 1 if and only if for all nonzero integers h we have

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N e^{2\pi i h x_n} = 0;$$

this is called **Weyl's criterion**. But if $x \in \Omega$, then

$$\left| \sum_{n=1}^N e^{2\pi i h n x} \right| = \left| \frac{1 - e^{2\pi i h N x}}{1 - e^{2\pi i h x}} \right| \leq \frac{2}{|1 - e^{2\pi i h x}|} = \frac{1}{|\sin \pi h x|}. \quad (19)$$

We thus obtain the following theorem.

Theorem 19. *If $x \in \Omega$ then the sequence $n x$ is uniformly distributed modulo 1.*

The **discrepancy** of a sequence ω is defined, for N a positive integer, by

$$D_N(\omega) = \sup_{0 \leq a < b \leq 1} \left| \frac{A([a, b]; N; \omega)}{N} - (b - a) \right|$$

One proves that the sequence ω is uniformly distributed modulo 1 if and only if $D_N(\omega) \rightarrow 0$ as $N \rightarrow \infty$ [85, p. 89, Theorem 1.1].

For $f : [0, 1] \rightarrow \mathbb{R}$, let $V(f)$ denote the total variation of f . Koksma's inequality [85, p. 143, Theorem 5.1] states that for any sequence $\omega = (x_n)$, for any $f : [0, 1] \rightarrow \mathbb{R}$ of bounded variation, and for any positive integer N , we have

$$\left| \frac{1}{N} \sum_{n=1}^N f(R(x_n)) - \int_0^1 f(t) dt \right| \leq V(f) D_N(\omega). \quad (20)$$

Following Kuipers and Niederreiter [85, p. 122, Lemma 3.2], we can bound the discrepancy of the sequence $n x$ in terms of the sum on the left-hand side of Theorem 14

Lemma 20. *There is some $C > 0$ such that for all $x \in \Omega$, $\omega = (nx)$, and for all positive integers m we have*

$$D_N(\omega) < C \left(\frac{1}{m} + \frac{1}{N} \sum_{j=1}^m \frac{1}{j \|jx\|} \right).$$

Proof. We shall use the following inequality, which lets us bound the discrepancy of a sequence in terms of exponential sums formed from the elements of the sequence. The Erdős-Turán theorem [85, p. 114, Eq. 2.42] states that there is some constant $C > 0$ such that for any sequence $\omega = (x_n)$ of real numbers, any positive integer N , and any positive integer m we have

$$D_N(\omega) \leq C \left(\frac{1}{m} + \sum_{j=1}^m \frac{1}{j} \left| \frac{1}{N} \sum_{n=1}^N e^{2\pi i j x_n} \right| \right). \quad (21)$$

Take $x_n = nx$. For each $j \geq 1$, by (19) we have

$$\left| \sum_{n=1}^N e^{2\pi i j n x} \right| \leq \frac{1}{|\sin \pi j x|} = \frac{1}{\sin(\pi \|jx\|)}.$$

But $\sin t \geq \frac{2}{\pi} t$ for $0 \leq t \leq \frac{\pi}{2}$, so

$$\frac{1}{\sin(\pi \|jx\|)} \leq \frac{1}{2 \|jx\|} < \frac{1}{\|jx\|}.$$

Using this in (21) gives us

$$D_N(\omega) < C \left(\frac{1}{m} + \sum_{j=1}^m \frac{1}{j} \cdot \frac{1}{N} \cdot \frac{1}{\|jx\|} \right) = C \left(\frac{1}{m} + \frac{1}{N} \sum_{j=1}^m \frac{1}{j \|jx\|} \right),$$

which is the claim. \square

It follows from Theorem 14 and Lemma 20 (taking $m = N$) that if x is of type $< K(\log h)^{1+\epsilon}$ then

$$D_N(\omega) = O \left(\frac{(\log N)^{2+\epsilon}}{N} \right). \quad (22)$$

Lemma 8 tells us that for almost all $x \in \Omega$ there is some $K > 0$ such that x is of type $< K(\log h)^{1+\epsilon}$, so for almost all $x \in \Omega$, the bound (22) is true. It likewise follows that if x has bounded partial quotients then

$$D_N(\omega) = O \left(\frac{(\log N)^2}{N} \right). \quad (23)$$

In fact, it can be proved that if $x \in \mathcal{B}_K$ then, for $g = \frac{1+\sqrt{5}}{2}$ [85, p. 125, Theorem 3.4],

$$D_N(\omega) \leq 3N^{-1} + \left(\frac{1}{\log g} + \frac{K}{\log(K+1)} \right) N^{-1} \log N.$$

We use the above bounds in the proof of the following theorem.

Theorem 21. *Let $\epsilon > 0$. For almost all x we have*

$$\sum_{n=1}^N \|nx\| = \frac{N}{4} + O((\log N)^{2+\epsilon}),$$

while if x has bounded partial quotients then

$$\sum_{n=1}^N \|nx\| = \frac{N}{4} + O((\log N)^2).$$

Proof. Let $f(t) = \|t\|$. Then $V(f) = 1$ and $\int_0^1 f(t)dt = \frac{1}{4}$, so we get from Koksma's inequality (20) that

$$\left| \frac{1}{N} \sum_{n=1}^N \|nx\| - \frac{1}{4} \right| \leq D_N(\omega),$$

thus

$$\sum_{n=1}^N \|nx\| = \frac{N}{4} + O(ND_N(\omega)).$$

The claims then follow respectively from (22) and (23). \square

Like we mentioned at the beginning of §6, because $|\sin(\pi x)| = \sin(\pi \|x\|) \leq \pi \|x\|$ and $|\sin(\pi x)| = \sin(\pi \|x\|) \geq \frac{2}{\pi} \cdot \pi \|x\| = 2 \|x\|$, we have

$$2 \sum_{n=1}^N \|nx\| \leq \sum_{n=1}^N |\sin(\pi nx)| \leq \pi \sum_{n=1}^N \|nx\|.$$

Thus Theorem 21 also gives estimates for $\sum_{n=1}^N |\sin(\pi nx)|$.

We can investigate the sum $\sum_{n=1}^N R(nx)$ rather than $\sum_{n=1}^N \|nx\|$; see Lang [92, p. 37, Theorem 1], who proves that for almost all $x \in \Omega$,

$$\sum_{n=1}^N R(nx) = \frac{N}{2} + O((\log N)^{2+\epsilon}).$$

For $x \in \Omega$, let $q_n = q_n(x)$, the denominator of the n th convergent of the continued fraction expansion of x , and let $a_n = a_n(x)$, the n th partial quotient

of the continued fraction expansion of x . For $m \geq 1$, one can prove [18, p. 211, Proposition 1] that m can be written in one and only one way in the form

$$m = \sum_{k=1}^{\infty} z_k q_{k-1} = \sum_{k=1}^t z_k q_{k-1}, \quad (24)$$

where (i) $0 \leq z_1 \leq a_1 - 1$, (ii) $0 \leq z_k \leq a_k$ for $k \geq 2$, (iii) for $k \geq 1$, if $z_{k+1} = a_{k+1}$ then $z_k = 0$, and (iv) $z_t \neq 0$ and $z_k = 0$ for $k > t$. The expression (24) is called the **Ostrowski expansion** of m . We emphasize that this expansion depends on x . Berthé [18] surveys applications of this numeration system in combinatorics. For $n \geq 0$, define $d_{2n} = q_{2n}x - p_{2n}$ and $d_{2n+1} = p_{2n+1} - q_{2n+1}x$. Brown and Shiue [24, p. 184, Theorem 1] prove that for $x \in \Omega$,

$$\sum_{k=1}^m \left(R(kx) - \frac{1}{2} \right) = \sum_{k=1}^t (-1)^k z_k \left(\frac{1}{2} - d_{k-1} \left(m_{k-1} + \frac{1}{2} z_k q_{k-1} + \frac{1}{2} \right) \right), \quad (25)$$

where $m_0 = 0$ and if $k \geq 1$ then $m_k = \sum_{j=1}^k z_j q_{j-1}$. If $k \geq 0$, then by (11) we have $0 < d_k < \frac{1}{q_{k+1}}$. For $k \geq 1$, using the fact that $q_k \geq m_k + 1$ (for the same reason that if the highest power of 2 appearing in a number's binary expansion is 2^{k-1} , then the number is $\leq 2^k - 1$),

$$\begin{aligned} m_{k-1} + \frac{1}{2} z_k q_{k-1} + \frac{1}{2} &= m_k - \frac{1}{2} z_k q_{k-1} + \frac{1}{2} \\ &\leq q_k - 1 - \frac{1}{2} z_k q_{k-1} + \frac{1}{2} \\ &= q_k - \frac{1}{2} - \frac{1}{2} z_k q_{k-1} \\ &< q_k. \end{aligned}$$

Using (25), this inequality, and the inequality $0 < d_k < \frac{1}{q_{k+1}}$, we obtain

$$\begin{aligned} \left| \sum_{k=1}^m \left(R(kx) - \frac{1}{2} \right) \right| &\leq \sum_{k=1}^t z_k \left| \frac{1}{2} - d_{k-1} \left(m_{k-1} + \frac{1}{2} z_k q_{k-1} + \frac{1}{2} \right) \right| \\ &< \frac{1}{2} \sum_{k=1}^t z_k \\ &< \frac{1}{2} \sum_{k=1}^t a_k. \end{aligned}$$

If the continued fraction expansion of x has bounded partial quotients, say $a_k \leq K$ for all k , we obtain from the above that

$$\left| \sum_{k=1}^m \left(R(kx) - \frac{1}{2} \right) \right| < \frac{Kt}{2}.$$

It can be proved [24, p. 185, Fact 2] that $t < 3 \log m$. Thus, if $a_k \leq K$ for all k , then for $m \geq 1$,

$$\left| \sum_{k=1}^m \left(R(kx) - \frac{1}{2} \right) \right| < \frac{3K \log m}{2}.$$

This is Lerch's claim stated in §3. For example, if $x = \frac{-1+\sqrt{5}}{2} \in \Omega$ then $a_k(x) = 1$ for all $k \geq 1$. We compute that

$$\sum_{k=1}^{1000000} \left(R(kx) - \frac{1}{2} \right) = 0.941799 \dots;$$

on the other hand, we compute that

$$\sum_{k=1}^{1000000} \left(R(k\pi) - \frac{1}{2} \right) = 19.223414 \dots$$

Brown and Shiue [24, p. 185, Fact 1] use (25) to obtain the result of Sierpinski stated in §3 that for all $x \in \Omega$,

$$\sum_{k=1}^m R(kx) = \frac{m}{2} + o(m).$$

They also prove [24, p. 188, Theorem 4] that for $A > 0$, there exists some $d_A > 0$ such that for $x \in \Omega$, if there are infinitely many t such that $\sum_{k=1}^t a_k \leq At$ (which happens in particular if x has bounded partial quotients), then there are infinitely many m such that

$$\sum_{k=1}^m \left(R(kx) - \frac{1}{2} \right) > d_A \log m,$$

and there are infinitely many m such that

$$\sum_{k=1}^m \left(R(kx) - \frac{1}{2} \right) < -d_A \log m.$$

It can be shown [92, p. 44, Theorem 4] that if k is a positive integer and $\epsilon > 0$, then for almost all x we have

$$\left| \sum_{n=1}^N e^{2\pi i n^k x} \right| = O\left(N^{\frac{1}{2} + \epsilon}\right).$$

Lang attributes this result to Vinogradov. But it is not so easy to obtain a bound on this exponential sum for specific x . For $k = 2$, one can prove [92, p. 45, Lemma] that for any $x \in \Omega$,

$$\left| \sum_{n=1}^N e^{2\pi i n^2 x} \right|^2 \leq N + 4 \sum_{n=1}^N \frac{1}{|\sin 4\pi n x|} < N + 4 \sum_{n=1}^{4N} \frac{1}{|\sin \pi n x|};$$

cf. Steele [143, Problem 14.2]. By (14) this gives us

$$\left| \sum_{n=1}^N e^{2\pi i n^2 x} \right|^2 < N + 2 \sum_{n=1}^{4N} \frac{1}{\|jx\|}.$$

If x has bounded partial quotients, it follows from Theorem 11 that

$$\left| \sum_{n=1}^N e^{2\pi i n^2 x} \right| = O\left(N^{1/2}(\log N)^{1/2}\right).$$

Hardy and Littlewood [59, p. 28, Theorem B5] prove that if $x \in \Omega$ is an algebraic number, then there is some $0 < \alpha(x) < 1$ such that $\sum_{n=1}^N R(nx) = \frac{N}{2} + O(N^{\alpha(x)})$. Pillai [115] gives a different proof of this.

Theorem 22 (Hardy and Littlewood, Pillai). *For $\tau > 2$, if $x \in \mathcal{D}(\tau)$ then for $\alpha = \frac{\tau-2}{\tau-1}$,*

$$\sum_{n=1}^N R(nx) = \frac{N}{2} + O(N^\alpha).$$

Pillai [114] proves other identities and inequalities for $\sum_{n=1}^N R(nx)$, some for all $x \in \Omega$ and some for all algebraic $x \in \Omega$.

For $\omega = (x_n)$, $n \geq 1$ and for $E \subset [0, 1)$, we remind ourselves that $A(E; M; \omega)$ denotes the number of x_n , $1 \leq n \leq M$, such that $R(x_n) \in E$. Define for $M \geq 1$,

$$D_M^*(\omega) = \sup_{0 < \beta \leq 1} \left| \frac{A([0, \beta); M; \omega)}{M} - \beta \right|.$$

It is straightforward to prove that $D_M^* \leq D_M \leq 2D_M^*$ [85, p. 91, Theorem 1.3]. For $N \geq 1$ write $r = \log N$. Let $\epsilon_0 > 0$ and let $N^{1/2} \leq \tau \leq N \exp(-r^{\epsilon_0})$. Suppose that $\alpha \in \mathbb{R}$ and

$$\alpha = \frac{a}{q} + \frac{\theta}{q\tau}, \quad \gcd(a, q) = 1, \quad \exp(r^{\epsilon_0}) \leq q \leq \tau, \quad |\theta| \leq 1.$$

For $0 < \beta < 1$ denote by $H_\beta(N)$ the number of primes $p \leq N$ such that $R(\alpha p) \leq \beta$. Vinogradov [150, p. 177, Chapter XI, Theorem] proves that for $\epsilon > 0$,

$$H_\beta(N) = \beta\pi(N) + O(N(q^{-1} + qN^{-1})^{\frac{1}{2}-\epsilon} + N^{\frac{4}{5}+\epsilon}), \quad N \rightarrow \infty.$$

Let $\omega = (p_n\alpha)$, $n \geq 1$, where p_n is the n th prime. Using Vinogradov's estimate, one proves that for $\alpha \in \mathbb{R} \setminus \mathbb{Q}$, $D_N^*(\omega) \rightarrow 0$ as $N \rightarrow \infty$, which implies that the sequence $(p_n\alpha)$ is uniformly distributed modulo 1. A clean proof of this is given by Pollicott [116, p. 200, Theorem 1], and this is also proved by Vaaler [146] using a Tauberian theorem. See also the early survey by Hua [68, pp. 98–99, §38].

Defining $S_\alpha(n) = \sum_{k=1}^n (R(k\alpha) - \frac{1}{2})$, Beck [10, p. 14, Theorem 3.1] proves that there is some $c > 0$ such that for every $\lambda \in \mathbb{R}$,

$$\frac{1}{N} \left| \left\{ 1 \leq n \leq N : \frac{S_{\sqrt{2}}(n)}{c\sqrt{\log n}} \leq \lambda \right\} \right| \rightarrow \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\lambda} \exp\left(-\frac{u^2}{2}\right) du$$

as $N \rightarrow \infty$. Beck [11, p. 20, Theorem 1.2] further proves that if α is a quadratic irrational, there are $C_1 = C_1(\alpha) \in \mathbb{R}$ and $C_2 = C_2(\alpha) \in \mathbb{R}_{>0}$ such that for $A, B \in \mathbb{R}$, $A < B$,

$$\begin{aligned} & \frac{1}{N} \left| \left\{ 0 \leq n < N : A \leq \frac{S_\alpha(n) - C_3 \log N}{C_4 \sqrt{\log N}} \leq B \right\} \right| \\ &= (2\pi)^{-1/2} \int_A^B e^{-u^2/2} du + O((\log N)^{-1/10} \log \log N). \end{aligned}$$

Beck and Chen [12]

9 Dirichlet series

The result of de la Vallée-Poussin [34] stated in §3 implies that there is no s such that for all irrational x the Dirichlet series $\sum_{n=1}^{\infty} \frac{n^{-s}}{|\sin n\pi x|}$ converges. It follows from this fact that there is no s such that for all irrational x the Dirichlet series $\sum_{n=1}^{\infty} \frac{n^{-s}}{\|nx\|}$ converges.

Lerch [97] in 1904 gives some statements without proof about the series

$$\sum_{\nu=1}^{\infty} \frac{\cot \nu\omega\pi}{(2\nu\pi)^{2m+1}}.$$

He states that if ω is a real algebraic number that does not belong to \mathbb{Q} , then for sufficiently large m this series converges, and states, for example, that with $\omega = \frac{1+\sqrt{5}}{2}$,

$$\sqrt{5} \sum_{\nu=1}^{\infty} \frac{\cot \nu\omega\pi}{(2\nu\pi)^7} = -\frac{8}{10!}.$$

Writing $c_r(\theta) = \sum_{n=1}^{\infty} \frac{\cot(\pi n\theta)}{n^{2r-1}}$, Berndt [17, p. 135, Theorem 5.1] proves that if θ is a real algebraic number of degree d and $1 < d < 2r - 1$ (to say that $d > 1$ is to say that θ is irrational), then $c_r(\theta)$ converges.

For a Dirichlet series $\sum_{n=1}^{\infty} a_n n^{-s}$, one can show [144, pp. 289–290, §9.11] that if the series is convergent at $s = \sigma_0 + it_0$, then for $\sigma > \sigma_0$ and any t the series is convergent at $s = \sigma + it$. It follows that there is some $\sigma_0 \in [-\infty, \infty]$ such if $\sigma < \sigma_0$ then the series diverges at $s = \sigma + it$, and if $\sigma > \sigma_0$ then the series converges at $s = \sigma + it$. We call σ_0 the **abscissa of convergence** of the Dirichlet series. If each a_n is a nonnegative real number, then the function

$$F(s) = \sum_{n=1}^{\infty} a_n n^{-s}, \quad \operatorname{Re} s > \sigma_0,$$

cannot be analytically continued to any domain that includes $s = \sigma_0$ [67, p. 101, Proposition 18].

Let a_n be a sequence of complex numbers. It can be shown [144, pp. 292–293, §9.14] that if $s_n = a_1 + \dots + a_n$ and the sequence s_n diverges, then the abscissa of convergence of the Dirichlet series $\sum_{n=1}^{\infty} a_n n^{-s}$ is given by

$$\sigma_0 = \limsup_{n \rightarrow \infty} \frac{\log |s_n|}{\log n}.$$

By Theorem 11 and Theorem 13 (taking, say, $\epsilon = 1$), for almost all $x \in \Omega$ there are C_1, C_2 such that for all positive integers n we have

$$C_1 n \log n < \sum_{j=1}^n \frac{1}{\|jx\|} < C_2 n (\log n)^2.$$

Thus, if $a_n = \frac{1}{\|nx\|}$, then

$$\frac{\log C_1}{\log n} + 1 + \frac{\log \log n}{\log n} < \frac{\log s_n}{\log n} < \frac{\log C_2}{\log n} + 1 + \frac{2 \log \log n}{\log n},$$

and hence

$$\lim_{n \rightarrow \infty} \frac{\log s_n}{\log n} = 1.$$

It follows that for almost all $x \in \Omega$ the abscissa of convergence of the Dirichlet series $\sum_{n=1}^{\infty} \frac{1}{\|nx\|} n^{-s}$ is $\sigma_0 = 1$.

Likewise, by Theorem 21 (taking $\epsilon = 1$), we get for almost all $x \in \Omega$ that $\sum_{j=1}^n \|jx\| = \frac{n}{4} + O((\log n)^3)$. We can then check that $\lim_{n \rightarrow \infty} \frac{\log s_n}{\log n} = 1$, and hence that the abscissa of convergence of the Dirichlet series $\sum_{n=1}^{\infty} \|nx\| n^{-s}$ is $\sigma_0 = 1$.

A 1953 result of Mahler [48, pp. 107–108] implies that if $\alpha \in \mathbb{R}$ is an algebraic number of degree d , then, for $m = \lfloor 20 \cdot 2^{\frac{5(d-1)}{2}} \rfloor$, the Dirichlet series

$$\sum_{n=1}^{\infty} \frac{n^{-s}}{\sin n\alpha}$$

has abscissa of convergence $\sigma_0 \leq d(m+1) \log(m+1)$, and the power series

$$\sum_{n=1}^{\infty} \frac{z^n}{\sin n\alpha}$$

has radius of convergence 1.

Rivoal [126] presents later work on similar Dirichlet series. See also Queffélec and Queffélec [120]. Lalín, Rodrigue and Rogers [88] prove results about Dirichlet series of the form $\sum_{n=1}^{\infty} \frac{n^{-s}}{\cos(n\pi z)}$. Duke and Imamoglu [42] review Hardy and Littlewood's work on estimating lattice points in triangles, and prove results about lattice points in cones.

For $\theta \in \Omega$, write

$$R_{r,\theta}(\zeta, m) = \sum_{n=0}^m \frac{1}{r^n} P_r(\zeta + n\theta),$$

where P_r is a periodic Bernoulli function. Spencer [142] proves that for any $\epsilon > 0$ and almost all $\theta \in \Omega$,

$$R_{1,\theta}(\zeta, m) = O((\log m)^{1+\epsilon}).$$

Another result Spencer proves in this paper is that if $q_n(\theta) = O(q_{n-1}^h)$, then $R_{r,\theta}(\zeta, m) = O(m^{1-\frac{r}{h}})$ for $1 \leq r < h$. Schoißengeier [133] gives an explicit formula for $\sum_{k=0}^{N-1} P_2(k\alpha)$.

10 Power series

For a power series $\sum a_n z^n$ with radius of convergence $0 \leq R \leq \infty$, the **Cauchy-Hadamard formula** [124, p. 111, Chapter 4, §1] states

$$R = \frac{1}{\limsup_{n \rightarrow \infty} |a_n|^{1/n}} = \liminf_{n \rightarrow \infty} |a_n|^{-1/n}. \quad (26)$$

The radius of convergence R is equal to the supremum of those $t \geq 0$ for which $|a_n|t^n$ is a bounded sequence.

Lemma 23. *If $x \in \Omega$, then the power series*

$$\sum_{n=1}^{\infty} \frac{z^n}{\|nx\|}$$

and

$$\sum_{n=1}^{\infty} \frac{z^n}{|\sin \pi nx|},$$

have the same radius of convergence.

Proof. The radii of convergence of these power series are respectively

$$\liminf_{n \rightarrow \infty} \|nx\|^{1/n} \quad \text{and} \quad \liminf_{n \rightarrow \infty} |\sin(\pi nx)|^{1/n}.$$

On the one hand,

$$|\sin(\pi nx)| = \sin(\pi \|nx\|) \leq \pi \|nx\|$$

Therefore, since $\lim_{n \rightarrow \infty} \pi^{1/n} = 1$,

$$\liminf_{n \rightarrow \infty} |\sin(\pi nx)|^{1/n} \leq \liminf_{n \rightarrow \infty} (\pi \|nx\|)^{1/n} = \liminf_{n \rightarrow \infty} \|nx\|^{1/n}$$

On the other hand, since $\sin t \geq \frac{2}{\pi}t$ for $t \geq 0$,

$$|\sin(\pi nx)| = \sin(\pi \|nx\|) \geq \frac{2}{\pi} \pi \|nx\| = 2 \|nx\|.$$

Therefore, using $\lim_{n \rightarrow \infty} 2^{1/n} = 1$, we have

$$\liminf_{n \rightarrow \infty} |\sin(\pi nx)|^{1/n} \geq \liminf_{n \rightarrow \infty} (2 \|nx\|)^{1/n} = \liminf_{n \rightarrow \infty} \|nx\|^{1/n},$$

showing that the two power series have the same radius of convergence. \square

We show in the following theorem that for almost all x , the power series $\sum_{n=1}^{\infty} \frac{z^n}{\|nx\|}$ has radius of convergence 1.

Theorem 24. *For almost all $x \in \Omega$, the power series*

$$\sum_{n=1}^{\infty} \frac{z^n}{\|nx\|} \tag{27}$$

has radius of convergence 1.

Proof. For $x \in \Omega$, let R_x be the radius of convergence of the power series (27). We have $0 < \|nx\| < \frac{1}{2}$, so $\frac{1}{\|nx\|} > 2$. Therefore $\sum_{n=1}^N \frac{1}{\|nx\|} \rightarrow \infty$ as $N \rightarrow \infty$, and so the power series (27) diverges at $z = 1$. Therefore $R_x \leq 1$ for all $x \in \Omega$.

We shall use Lemma 7 to get a lower bound on R_x that holds for almost all $x \in \Omega$. Let $A = \{x \in \Omega : R_x < 1\}$, let $A_m = \{x \in \Omega : R_x < 1 - \frac{1}{m}\}$, and let B_m be those $x \in \Omega$ such that $\|nx\|^{1/n} < 1 - \frac{1}{m}$ infinitely often. If $x \in A_m$, then $R_x = \liminf_{n \rightarrow \infty} \|nx\|^{1/n} < 1 - \frac{1}{m}$, and this implies that there are infinitely many n such that $\|nx\|^{1/n} < 1 - \frac{1}{m}$, so $x \in B_m$, i.e. $A_m \subset B_m$. But let $f_m(n) = (1 - \frac{1}{m})^n$. Then $\sum_{n=1}^{\infty} f_m(n)$ converges, since $1 - \frac{1}{m} < 1$, so, by Lemma 7, for almost all $x \in \Omega$ there are only finitely many n such that $\|nx\| < f_m(n)$. Thus $\mu(B_m) = 0$. Hence $\mu(A_m) = 0$, and since $A = \bigcup_{m=2}^{\infty} A_m$ we get that

$$\mu(A) \leq \sum_{m=2}^{\infty} \mu(A_m) = 0,$$

that is, $R_x \geq 1$ for almost all $x \in \Omega$. In conclusion, $R_x = 1$ for almost all $x \in \Omega$. \square

In fact, we can prove the above theorem using the bounds we obtained in Theorem 11. By Theorem 11, for almost all $x \in \Omega$ we have that $\sum_{j=1}^m \frac{1}{\|jx\|} = O(m^2)$. (Here we will merely need the fact that the sum is subexponential in m .) For such an x , take $0 < r < 1$. Let $a_n = r^n$, let $s_n = \sum_{j=1}^n \frac{1}{\|jx\|}$, let $b_1 = 0$, and let $b_n = s_{n-1}$ for $n \geq 2$. Using summation by parts, namely

$$\sum_{n=1}^N a_n (b_{n+1} - b_n) = a_{N+1} b_{N+1} - a_1 b_1 - \sum_{n=1}^N b_{n+1} (a_{n+1} - a_n),$$

we get

$$\sum_{n=1}^N \frac{r^n}{\|nx\|} = r^{N+1} s_N + \sum_{n=1}^N s_n r^n (1-r).$$

Therefore

$$\sum_{n=1}^N \frac{r^n}{\|nx\|} = O(r^{N+1} N^2) + O\left(\sum_{n=1}^N n^2 r^n\right) = O(1).$$

Since $\sum_{n=1}^N \frac{r^n}{\|nx\|}$ is increasing in N (being a sum of positive terms), we obtain that the series $\sum_{n=1}^{\infty} \frac{r^n}{\|nx\|}$ converges. Since this is true for all r with $0 < r < 1$, it follows that $R_x \geq 1$.

For $x \in \Omega$ let R_x be the radius of convergence of the power series $\sum_{q=1}^{\infty} \frac{z^q}{\|qx\|}$. We proved in Theorem 27 that for almost all $x \in \Omega$, $R_x = 1$.

Theorem 25. *For $x \in \Omega$, let R_x be the radius of convergence of the power series $\sum_{q=1}^{\infty} \frac{z^q}{\|qx\|}$, and let $a_n = a_n(x)$ and $q_n = q_n(x)$. Then*

$$R_x = \liminf_{n \rightarrow \infty} a_{n+1}^{-1/q_n}.$$

For any $0 \leq R \leq 1$ there is some $x \in \Omega$ such that $R_x = R$.

Proof. From the Cauchy-Hadamard formula (26),

$$R_x = \liminf_{q \rightarrow \infty} \|qx\|^{1/q}.$$

Then $R_x \leq \liminf_{n \rightarrow \infty} \|q_n x\|^{1/q_n}$. On the one hand, by (11), $\|q_n x\| < q_{n+1}^{-1}$, and $q_{n+1} = a_{n+1} q_n + q_{n-1} > a_{n+1} q_n$ hence $\|q_n x\| < a_{n+1}^{-1} q_n^{-1}$, and using that $\lim_{n \rightarrow \infty} q_n^{-1/q_n} = 1$,

$$R_x \leq \liminf_{n \rightarrow \infty} a_{n+1}^{-1/q_n} q_n^{-1/q_n} = \liminf_{n \rightarrow \infty} a_{n+1}^{-1/q_n}.$$

On the other hand, let $q \geq 2$ and take $q_n \leq q < q_{n+1}$. Applying (12),

$$\|q_n x\| > \frac{1}{q_{n+1} + q_n} > \frac{1}{2q_{n+1}} = \frac{1}{2(a_{n+1} q_n + q_{n-1})} > \frac{1}{4a_{n+1} q_n}.$$

Then applying Theorem 2, and using that $0 < \|q_n x\| < 1$ and $q \geq q_n$,

$$\|qx\|^{1/q} \geq \|q_n x\|^{1/q} \geq \|q_n x\|^{1/q_n} > \left(\frac{1}{4a_{n+1} q_n}\right)^{1/q_n}.$$

As $(4q_n)^{-1/q_n} \rightarrow 1$ as $n \rightarrow \infty$, this implies

$$R_x = \liminf_{q \rightarrow \infty} \|qx\|^{1/q} \geq \liminf_{n \rightarrow \infty} a_{n+1}^{-1/q_n}.$$

For $0 < R < 1$, let $R = e^{-r}$ for $r > 0$. Define $a \in \mathbb{N}^{\mathbb{N}}$ as follows. Define $a_1 = 1$. Suppose for $n \geq 1$ that we have defined a_1, \dots, a_n and thus p_1, \dots, p_n

and q_1, \dots, q_n . Define $a_{n+1} = [e^{r q_n}]$. Then $a_{n+1}^{1/q_n} \leq e^r$, so $a_{n+1}^{-1/q_n} \geq e^{-r}$. Therefore for $x = v(a)$, $R_x \geq e^{-r}$. Now, $e^{r q_n} > 1$ so $a_{n+1} = [e^{r q_n}] \geq 2^{-1} e^{r q_n}$. Then $a_{n+1}^{1/q_n} \geq 2^{-1/q_n} e^r$, hence

$$R_x = \liminf_{n \rightarrow \infty} a_{n+1}^{-1/q_n} \leq \liminf_{n \rightarrow \infty} 2^{1/q_n} e^{-r} = e^{-r}.$$

We have therefore established that when $x = v(a)$, $R_x = e^{-r}$.

For $R = 0$, define $a \in \mathbb{N}^{\mathbb{N}}$ by $a_1 = 1$ and $a_{n+1} = [e^{n q_n}]$, which satisfies $a_{n+1} \geq 2^{-1} e^{n q_n}$. For $x = v(a)$,

$$R_x = \liminf_{n \rightarrow \infty} a_{n+1}^{-1/q_n} \leq \liminf_{n \rightarrow \infty} 2^{1/q_n} e^{-n} = 0.$$

For $R = 1$, define $a \in \mathbb{N}^{\mathbb{N}}$ by $a_n = 1$ for all $n \geq 1$. Namely, $v(a) = \frac{-1+\sqrt{5}}{2} \in \Omega$. For $x = v(a)$ it is immediate that $R_x \geq 1$. \square

Since $R(nx) < 1$, of course the power series $\sum_{n=1}^{\infty} R(nx)z^n$ has radius of convergence ≥ 1 . The following result, for which Pólya and Szegő [117, p. 280, Part II, No. 168] cite Hecke, shows in particular that the radius of convergence of this power series is ≤ 1 for $x \in \Omega$ and is thus equal to 1.

Theorem 26. For $x \in \Omega$, let

$$f(z) = \sum_{n=1}^{\infty} R(nx)z^n, \quad |z| < 1.$$

We have

$$\lim_{r \rightarrow 1^-} (1-r)f(re^{2\pi i x}) = \frac{1}{2\pi i}.$$

Proof. Since $x \in \Omega$, the sequence nx is uniformly distributed modulo 1. Therefore, with $f(t) = te^{2\pi i t}$ we have by (18) that

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N R(nx)e^{2\pi i n x} = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N f(R(nx)) = \int_0^1 te^{2\pi i t} dt = \frac{1}{2\pi i}.$$

We will use the following result [117, p. 21, Part I, No. 88]. If a sequence of complex numbers a_n satisfies $\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N a_n = s$, then

$$\lim_{t \rightarrow 1^-} (1-t) \sum_{n=1}^{\infty} a_n t^n = s.$$

Let $a_n = R(nx)e^{2\pi i n x}$, and we thus have

$$\lim_{t \rightarrow 1^-} (1-t) \sum_{n=1}^{\infty} R(nx)e^{2\pi i n x} t^n = \frac{1}{2\pi i}.$$

\square

It follows from the above theorem that if $x \in \Omega$ then $|z| = 1$ is a natural boundary of the function f defined on the open unit disc by $f(z) = \sum_{n=1}^{\infty} R(nx)z^n$; cf. Segal [134, p. 255, Chapter 6], who writes about this power series, and who gives a thorough introduction to natural boundaries in the same chapter. Breur and Simon [22] prove a generalization of this result.

Hata [62, p. 173, Problem 12.6] mentions the appearance of the function f from the above theorem in the study of the Caianiello neuron equations.

11 Product

We will use the following lemma proved by Hardy and Littlewood [60, p. 89], whose brief proof we expand.

Lemma 27. *Let $\psi : (0, \infty) \rightarrow \mathbb{R}$ be positive and nondecreasing. If*

$$\sum_{k=1}^{\infty} \frac{1}{k\psi(k)} < \infty,$$

then for almost all $x \in \Omega$, there exists some H such that for all $n \geq 1$ and for all real $h \geq H$, there are at most $\max\{\frac{n\psi(h)}{h}, 1\}$ integers $m \in \{1, \dots, n\}$ that satisfy $\|mx\| < \frac{1}{h}$.

Proof. By Lemma 7, for almost all $x \in \Omega$ there is some K such that if $k \geq K$ then

$$\|kx\| \geq \frac{2}{k\psi(k)}. \quad (28)$$

Let H be large enough so that

$$\min_{k < K} \|kx\| \geq \frac{2}{H};$$

also let $\psi(H) \geq 1$. Now suppose by contradiction that there is some $n \geq 1$ and some $h \geq H$ such that there are more than $\max\{\frac{n\psi(h)}{h}, 1\}$ integers $m \in \{1, \dots, n\}$ that satisfy $\|mx\| < \frac{1}{h}$. Then there are some $1 \leq m_1 < m_2 \leq n$ satisfying $\|m_1x\| < \frac{1}{h}$ and $\|m_2x\| < \frac{1}{h}$ and such that

$$\mu = m_2 - m_1 < \frac{n}{\frac{n\psi(h)}{h}} = \frac{h}{\psi(h)},$$

so $\mu\psi(h) < h$. On the other hand,

$$\|\mu x\| \leq \|m_1x\| + \|m_2x\| < \frac{1}{h} + \frac{1}{h} = \frac{2}{h},$$

so $h < \frac{2}{\|\mu x\|}$. Thus $\mu\psi(h) < \frac{2}{\|\mu x\|}$, i.e.,

$$\|\mu x\| < \frac{2}{\mu\psi(h)} \leq \frac{2}{\mu\psi(\mu)};$$

$\mu < h$ because $\mu < \frac{h}{\psi(h)}$ and $\psi(h) \geq \psi(H) \geq 1$. Moreover, since $\|\mu x\| < \frac{2}{h} \leq \frac{2}{H}$, we have $\mu \geq K$. This contradicts (28). \square

Hardy and Littlewood [60, p. 89, Theorem 4] prove the following theorem that gives us the conclusion (18) for certain functions that are not Riemann integrable on $[0, 1]$.

Theorem 28. *Let $f : (0, 1) \rightarrow \mathbb{R}$ be nonnegative, let f be nonincreasing on $(0, \frac{1}{2})$ and nondecreasing on $(\frac{1}{2}, 1)$, and let*

$$\int_0^1 f(t)dt < \infty.$$

Let $\psi : (1, \infty) \rightarrow \mathbb{R}$ be a positive and nondecreasing function such that

$$\sum_{k=2}^{\infty} \frac{1}{k\psi(k)} < \infty.$$

If

$$\int_0^1 f(t) \left(\psi\left(\frac{1}{t}\right) + \psi\left(\frac{1}{1-t}\right) \right) dt < \infty,$$

then for almost all $x \in \Omega$,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{m=1}^n f(R(mx)) = \int_0^1 f(t)dt.$$

Proof. For $0 < \delta < \frac{1}{2}$, define

$$f_{\delta}(t) = \begin{cases} f(t), & \delta \leq t \leq 1 - \delta, \\ 0, & 0 < t < \delta \text{ or } 1 - \delta < t < 1. \end{cases}$$

From Lemma 27, for almost all $x \in \Omega$ there is some C such that for all n and h there are at most $C \frac{n\psi(h)}{h}$ integers $m \in \{1, \dots, n\}$ satisfying $\|mx\| < \frac{1}{h}$. Let

$$S_n = \frac{1}{n} \sum_{m=1}^n f(R(mx)) = S_n^1(\delta) + S_n^2(\delta),$$

where

$$S_n^1(\delta) = \frac{1}{n} \sum_{m=1}^n f_{\delta}(R(mx))$$

and

$$\begin{aligned}
S_n^2(\delta) &= \frac{1}{n} \sum_{m=1}^n f(R(mx)) - f_\delta(R(mx)) \\
&= \frac{1}{n} \sum_{\substack{1 \leq m \leq n \\ \|mx\| < \delta}} f(R(mx)) \\
&= \frac{1}{n} \sum_{k=0}^{\infty} \sum_{\substack{1 \leq m \leq n \\ \frac{\delta}{2^{k+1}} \leq \|mx\| < \frac{\delta}{2^k}}} f(R(mx)) \\
&= \frac{1}{n} \sum_{k=0}^{\infty} T_{k,n}(\delta).
\end{aligned}$$

There are at most $C \frac{n\psi\left(\frac{2^k}{\delta}\right)}{2^k}$ integers $m \in \{1, \dots, n\}$ that satisfy $\|mx\| < \frac{\delta}{2^k}$,

thus, as ψ is nondecreasing, there are at most $2C \frac{n\delta\psi\left(\frac{2^k}{\delta}\right)}{2^{k+1}} \leq 2C \frac{n\delta\psi\left(\frac{2^{k+1}}{\delta}\right)}{2^{k+1}}$ terms in $T_{k,n}(\delta)$. For each term $f(R(mx))$ in $T_{k,n}(\delta)$, since $\frac{\delta}{2^{k+1}} \leq \|mx\|$ we have, by assumption on f , either $f(R(mx)) \leq f\left(\frac{\delta}{2^{k+1}}\right)$ or $f(R(mx)) \leq f\left(1 - \frac{\delta}{2^{k+1}}\right)$, and hence

$$f(R(mx)) \leq f\left(\frac{\delta}{2^{k+1}}\right) + f\left(1 - \frac{\delta}{2^{k+1}}\right).$$

Therefore,

$$\begin{aligned}
S_n^2(\delta) &\leq \frac{1}{n} \sum_{k=0}^{\infty} 2C \frac{n\delta\psi\left(\frac{2^{k+1}}{\delta}\right)}{2^{k+1}} \left(f\left(\frac{\delta}{2^{k+1}}\right) + f\left(1 - \frac{\delta}{2^{k+1}}\right) \right) \\
&= 4C \sum_{k=0}^{\infty} \frac{\delta}{2^{k+2}} \psi\left(\frac{2^{k+1}}{\delta}\right) f\left(\frac{\delta}{2^{k+1}}\right) + 4C \sum_{k=0}^{\infty} \frac{\delta}{2^{k+2}} \psi\left(\frac{2^{k+1}}{\delta}\right) f\left(1 - \frac{\delta}{2^{k+1}}\right) \\
&\leq 4C \int_0^{\frac{\delta}{2}} \psi\left(\frac{1}{t}\right) f(t) dt + 4C \int_{1-\frac{\delta}{2}}^1 \psi\left(\frac{1}{1-t}\right) f(t) dt.
\end{aligned}$$

Let $\epsilon > 0$. Because $\int_0^1 f(t) \left(\psi\left(\frac{1}{t}\right) + \psi\left(\frac{1}{1-t}\right) \right) dt < \infty$, there exists a δ_1 such that if $\delta \leq \delta_1$ then $S_n^2(\delta) < \epsilon$.

On the other hand, since f_δ is Riemann integrable on $[0, 1]$ and because, by (19), the sequence mx is uniformly distributed modulo 1, we obtain from (18) that

$$\lim_{n \rightarrow \infty} S_n^1(\delta) = \int_0^1 f_\delta(t) dt = \int_\delta^{1-\delta} f(t) dt.$$

As $\int_0^1 f(t) dt < \infty$, there exists a δ_2 such that for $\delta \leq \delta_2$ and for sufficiently large n ,

$$\left| S_n^1(\delta) - \int_0^1 f(t) dt \right| \leq \left| S_n^1(\delta) - \int_\delta^{1-\delta} f(t) dt \right| + \left| \int_0^\delta f(t) dt \right| + \left| \int_{1-\delta}^1 f(t) dt \right| < 3\epsilon.$$

Therefore, for sufficiently large n and for sufficiently small δ ,

$$\left| S_n - \int_0^1 f(t) dt \right| \leq \left| S_n^1(\delta) - \int_0^1 f(t) dt \right| + |S_n^2(\delta)| < 4\epsilon.$$

Thus for sufficiently large n ,

$$\left| S_n - \int_0^1 f(t) dt \right| \leq 4\epsilon.$$

□

By the Birkhoff ergodic theorem [44, p. 44, Theorem 2.30], if $f \in L^1[0, 1]$ and $x \in \Omega$, then for almost all $\alpha \in [0, 1]$,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=1}^n f(R(\alpha + jx)) = \int_0^1 f(t) dt.$$

This equality holding for $\alpha = 0$ is the conclusion of Theorem 28.

Baxa [9] reviews further results that give conditions when a function $f : [0, 1] \rightarrow \mathbb{R} \cup \{+\infty\}$ that is not Riemann integrable on $[0, 1]$ nevertheless satisfies

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{m=1}^n f(R(mx)) = \int_0^1 f(t) dt$$

for certain $x \in \Omega$. Oskolkov [109, p. 170, Theorem 1] shows that if $f : (0, 1) \rightarrow \mathbb{R}$ satisfies $\lim_{t \rightarrow 0+} f(t) = +\infty$ and $\lim_{t \rightarrow 1-} f(t) = +\infty$, and also the improper Riemann integral of f on $[0, 1]$ exists, then, for $x \in \Omega$,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{m=1}^n f(R(mx)) = \int_0^1 f(t) dt$$

if and only if

$$\lim_{n \rightarrow \infty} \frac{1}{q_n(x)} f(R(q_n(x)x)) = 0,$$

where $q_n(x)$ is the denominator of the n th convergent of the continued fraction expansion of x .

Driver, Lubinsky, Petruska and Sarnak [40]

Using Theorem 28 we can now prove the following theorem of Hardy and Littlewood [60, p. 88, Theorem 2].

Theorem 29. *For almost all $x \in \Omega$,*

$$\lim_{n \rightarrow \infty} \left(\prod_{k=1}^n |\sin k\pi x| \right)^{1/n} = \frac{1}{2}.$$

Proof. Let $f(t) = -\log \sin \pi t$. Using $\cos(t - \frac{\pi}{2}) = \sin t$ and $\sin 2t = 2 \sin t \cos t$, one can check that $\int_0^1 \log \sin \pi t dt = -\log 2$. (The earliest evaluation of this integral of which we are aware is by Euler [46], who gives two derivations, the first using the Euler-Maclaurin summation formula, the power series expansion for $\log\left(\frac{1+z}{1-z}\right)$, and the power series expansion of $z \cot(z)$, and the second using the Fourier series of $\log |\sin t|$.) Thus, $\int_0^1 f(t) dt = \log 2 < \infty$. So f satisfies the conditions of Theorem 28.

Let $\psi(t) = (\log t)^2$. First, upper bounding the series by an integral,

$$\sum_{k=2}^{\infty} \frac{1}{k(\log k)^2} \leq \frac{1}{2(\log 2)^2} + \int_2^{\infty} \frac{1}{t(\log t)^2} dt = \frac{1}{2(\log 2)^2} + \log 2 < \infty.$$

Second,

$$\begin{aligned} \int_0^1 f(t) \left(\psi\left(\frac{1}{t}\right) + \psi\left(\frac{1}{1-t}\right) \right) dt &= \int_0^1 f(t) (\psi(t) + \psi(1-t)) dt \\ &= \int_0^{\frac{1}{2}} -2 \log \sin(\pi t) \\ &\quad ((\log t)^2 + (\log(1-t))^2) dt \\ &\leq \int_0^{\frac{1}{2}} -2 \log(2t) ((\log t)^2 + (\log(1-t))^2) dt \\ &< \infty. \end{aligned}$$

Therefore by Theorem 28, for almost all $x \in \Omega$,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{m=1}^n -\log \sin(\pi R(mx)) = \int_0^1 -\log \sin \pi t dt,$$

i.e.

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{m=1}^n \log |\sin \pi mx| = \log \frac{1}{2}.$$

□

Hardy and Littlewood give another proof [60, p. 86, Theorem 1] of the above theorem, which we now work out. This proof is complicated and we greatly expand on the abbreviated presentation of Hardy and Littlewood.

We remind ourselves that the Cauchy-Hadamard formula states that the radius of convergence R of a power series $\sum a_n z^n$ satisfies

$$R = \frac{1}{\limsup_{n \rightarrow \infty} |a_n|^{1/n}} = \liminf_{n \rightarrow \infty} |a_n|^{-1/n}.$$

Theorem 30. Fix $x \in \Omega$ and write $q_0 = e^{2\pi i x}$. Let ρ and R respectively be the radii of convergence of the power series

$$f(z) = \sum_{n=1}^{\infty} \frac{z^n}{n(1 - q_0^n)}, \quad F(z) = 1 + \sum_{n=1}^{\infty} \frac{z^n}{(1 - q_0)(1 - q_0^2) \cdots (1 - q_0^n)}.$$

Then $R = \rho$, and if $|z| < \rho$ then $F(z) = e^{f(z)}$.

The functions $f : D(0, \rho) \rightarrow \mathbb{C}$ and $F : D(0, R) \rightarrow \mathbb{C}$ are analytic [144, p. 69, Theorem 2.16].

Using the Cauchy-Hadamard formula we have

$$\rho = \liminf_{n \rightarrow \infty} n^{1/n} |1 - q_0^n|^{1/n} \leq \liminf_{n \rightarrow \infty} (2n)^{1/n} = 1$$

and

$$R = \liminf_{n \rightarrow \infty} (|1 - q_0| |1 - q_0^2| \cdots |1 - q_0^n|)^{1/n} \leq 2\rho. \quad (29)$$

Lemma 31. For $|u| = 1$ and for $0 \leq r < 1$,

$$\left| \frac{1 - u}{1 - ru} \right| \leq 2.$$

Proof.

$$|1 - u| \leq |1 - ru| + |ru - u| = |1 - ru| + 1 - r.$$

Because $\operatorname{Re} u \leq 1$ we have $1 - r \leq 1 - r\operatorname{Re} u = \operatorname{Re}(1 - ru) \leq |1 - ru|$. Hence $|1 - u| \leq 2|1 - ru|$, from which the claim follows. \square

We assert the following as a common fact in complex analysis.

Lemma 32. For $|w| < 1$ define

$$L(w) = \sum_{n=1}^{\infty} \frac{w^n}{n}.$$

If $|w| < 1$, then $e^{L(w)} = (1 - w)^{-1}$.

For $|q|, |z| < 1$, we define

$$f(z, q) = \sum_{n=1}^{\infty} \frac{z^n}{n(1 - q^n)}, \quad F(z, q) = 1 + \sum_{n=1}^{\infty} \frac{z^n}{(1 - q)(1 - q^2) \cdots (1 - q^n)}.$$

For $|q| < 1$, define $c_0(q) = 0$ and for $n \geq 1$,

$$c_n(q) = \frac{1}{n(1 - q^n)},$$

and thus for $|z| < 1$,

$$f(z, q) = \sum_{n=0}^{\infty} c_n(q) z^n.$$

Furthermore define $\gamma_0 = 0$ and for $n \geq 1$,

$$\gamma_n = \frac{1}{n(1 - q_0^n)},$$

and thus for $|z| < \rho$,

$$f(z) = \sum_{n=0}^{\infty} \gamma_n z^n.$$

For $|q| < 1$, define $C_0(q) = 1$ and for $n \geq 1$,

$$C_n(q) = \frac{1}{(1-q)(1-q^2)\cdots(1-q^n)},$$

and thus for $|z| < 1$,

$$F(z, q) = \sum_{n=0}^{\infty} C_n(q) z^n.$$

Furthermore define $\Gamma_0 = 1$ and for $n \geq 1$,

$$\Gamma_n = \frac{1}{(1-q_0)(1-q_0^2)\cdots(1-q_0^n)},$$

and thus for $|z| < R$,

$$F(z) = \sum_{n=0}^{\infty} \Gamma_n z^n.$$

We prove directly the following, which is an instance of the **q -binomial formula** [3, p. 17, Theorem 2.1].

Proposition 33. *If $|q|, |z| < 1$ then*

$$F(z, q) = e^{f(z, q)}.$$

Proof. By Lemma 32,

$$f(z, q) = \sum_{n=1}^{\infty} \frac{z^n}{n} \left(\sum_{m=0}^{\infty} q^{nm} \right) = \sum_{m=0}^{\infty} \left(\sum_{n=1}^{\infty} \frac{(zq^m)^n}{n} \right) = \sum_{m=0}^{\infty} L(zq^m).$$

Because $e^{L(w)} = (1-w)^{-1}$ for $|w| < 1$,

$$e^{f(z, q)} = \prod_{m=0}^{\infty} e^{L(zq^m)} = \prod_{m=0}^{\infty} (1 - zq^m)^{-1}.$$

Define

$$G(z, q) = \prod_{m=0}^{\infty} (1 - zq^m)^{-1} = \sum_{n=0}^{\infty} g_n(q) z^n.$$

On the one hand, $g_0(q) = G(0, q) = 1$. On the other hand,

$$G(qz, q) = (1-z)G(z, q),$$

thus

$$\sum_{n=0}^{\infty} g_n(q)z^{n+1} = \sum_{n=0}^{\infty} g_n(q)(1-q^n)z^n,$$

and therefore for $n \geq 1$ we have $g_n(q) = (1-q^n)^{-1}g_{n-1}(q)$. Thus by induction, for $n \geq 1$,

$$g_n(q) = \frac{1}{(1-q)(1-q^2)\cdots(1-q^n)}.$$

Hence

$$e^{f(z,q)} = 1 + \sum_{n=1}^{\infty} \frac{z^n}{(1-q)(1-q^2)\cdots(1-q^n)} = F(z,q).$$

□

Proposition 34. $R \geq \rho$, and if $|z| < \rho$ then $e^{f(z)} = F(z)$.

Proof. If $\rho = 0$ then the claim is immediate. Otherwise, $0 < \rho \leq 1$. Let $0 < t < \rho$, $0 < r < 1$, and define $G(\theta) = F(te^{i\theta}, rq_0)$. On the one hand,

$$G(\theta) = \sum_{n=0}^{\infty} C_n(rq_0)(te^{i\theta})^n = \sum_{n=0}^{\infty} C_n(rq_0)t^n e^{in\theta},$$

which implies that $\widehat{G}(n) = C_n(rq_0)t^n$ for $n \geq 0$ and $\widehat{G}(n) = 0$ for $n < 0$. On the other hand, for $n \in \mathbb{Z}$, using Proposition 33,

$$\begin{aligned} \widehat{G}(n) &= \frac{1}{2\pi} \int_0^{2\pi} G(\theta)e^{-in\theta} d\theta \\ &= \frac{1}{2\pi} \int_0^{2\pi} F(te^{i\theta}, rq_0)e^{-in\theta} d\theta \\ &= \frac{1}{2\pi} \int_0^{2\pi} e^{f(te^{i\theta}, rq_0)} e^{-in\theta} d\theta. \end{aligned}$$

By Lemma 31,

$$|f(te^{i\theta}, rq_0)| \leq \sum_{n=1}^{\infty} \frac{t^n}{n|1-r^nq_0^n|} \leq 2 \sum_{n=1}^{\infty} \frac{t^n}{n|1-q_0^n|} = M(t),$$

and because $t < \rho$ it is the case that $M(t) < \infty$. Then for $n \geq 0$,

$$|C_n(rq_0)t^n| = |\widehat{G}(n)| \leq \frac{1}{2\pi} \int_0^{2\pi} |e^{f(te^{i\theta}, rq_0)}| d\theta \leq e^{M(t)}.$$

$C_n(rq_0) \rightarrow \Gamma_n$ as $r \rightarrow 1$, and therefore

$$|\Gamma_n t^n| \leq e^{M(t)}, \quad 0 < t < \rho, \quad n \geq 0.$$

Now fix $|z| < \rho$, take t such that $|z| < t < \rho$, and write $0 \leq \delta = \frac{|z|}{t} < 1$ and $M_n = e^{M(t)}\delta^n$. Because $|\Gamma_n t^n| \leq e^{M(t)}$,

$$\sum_{n=0}^{\infty} |\Gamma_n z^n| = \sum_{n=0}^{\infty} \delta^n |\Gamma_n t^n| \leq \sum_{n=0}^{\infty} M_n = \frac{e^{M(t)}}{1-\delta} < \infty,$$

which implies that $|z| \leq R$. Therefore $R \geq \rho$. Furthermore, because

$$|C_n(rq_0)z^n| = \delta^n |C_n(rq_0)t^n| = M_n, \quad r \in (0, 1),$$

by the Weierstrass M -test [129, p. 148, Theorem 7.10], the sequence $\sum_{n=0}^N C_n(rq_0)z^n$ converges uniformly for $r \in (0, 1)$ and therefore [129, p. 149, Theorem 7.11]

$$\begin{aligned} \lim_{r \rightarrow 1} F(z, rq_0) &= \lim_{r \rightarrow 1} \lim_{N \rightarrow \infty} \sum_{n=0}^N C_n(rq_0)z^n \\ &= \lim_{N \rightarrow \infty} \lim_{r \rightarrow 1} \sum_{n=0}^N C_n(rq_0)z^n \\ &= \lim_{N \rightarrow \infty} \sum_{n=0}^N \Gamma_n z^n \\ &= F(z). \end{aligned}$$

Now, for $0 < r < 1$, by Lemma 31,

$$|c_n(rq_0)z^n| = \left| \frac{z^n}{n(1-r^n q_0^n)} \right| \leq 2 \left| \frac{z^n}{n(1-q_0^n)} \right| = m_n.$$

Because $|z| < \rho$, the series $\sum_{n=0}^{\infty} m_n$ converges, and therefore by the Weierstrass M -test, the sequence $\sum_{n=0}^N c_n(rq_0)z^n$ converges uniformly for $r \in (0, 1)$. Then

$$\begin{aligned} \lim_{r \rightarrow 1} f(z, rq_0) &= \lim_{r \rightarrow 1} \lim_{N \rightarrow \infty} \sum_{n=0}^N c_n(rq_0)z^n \\ &= \lim_{N \rightarrow \infty} \lim_{r \rightarrow 1} \sum_{n=0}^N c_n(rq_0)z^n \\ &= \lim_{N \rightarrow \infty} \sum_{n=0}^N \gamma_n z^n \\ &= f(z). \end{aligned}$$

Then using Lemma 33,

$$\begin{aligned} \exp(f(z)) &= \exp\left(\lim_{r \rightarrow 1} f(z, rq_0)\right) \\ &= \lim_{r \rightarrow 1} \exp(f(z, rq_0)) \\ &= \lim_{r \rightarrow 1} F(z, rq_0) \\ &= F(z), \end{aligned}$$

completing the proof. \square

Lemma 35. *If $|u| = 1$ and $u \neq 1$ then*

$$\left| \sum_{n=1}^N u^n \right| \leq \frac{2}{|1-u|}.$$

Proof.

$$\left| \sum_{n=1}^N u^n \right| = \left| \frac{u - u^{N+1}}{1-u} \right| = \left| \frac{1 - u^N}{1-u} \right| \leq \frac{2}{|1-u|}.$$

\square

Lemma 36. *Let $0 < a < 1$, let $x \in \Omega$, and suppose that there is some C such that $\frac{a^n}{|\sin n\pi x|} \leq C$ for all n . Then*

$$\sum_{n=1}^N \frac{a^{2n}}{\sin^2 n\pi x} = o(N), \quad N \rightarrow \infty.$$

Proof. Take $0 < \epsilon < \frac{1}{2}$ and let

$$E_N = \{n : 1 \leq n \leq N, \|nx\| < \epsilon\}.$$

Thus if $1 \leq n \leq N$ and $n \notin E_N$ then $\|nx\| \geq \epsilon$. Write $S_N = \sum_{n=1}^N \frac{a^{2n}}{\sin^2 n\pi x}$. Because $|\sin n\pi x| = \sin(\pi \|nx\|) \geq 2\|nx\|$,

$$\begin{aligned} S_N &= \sum_{n \in E_N} \frac{a^{2n}}{\sin^2 n\pi x} + \sum_{n \notin E_N, 1 \leq n \leq N} \frac{a^{2n}}{\sin^2 n\pi x} \\ &\leq C^2 |E_N| + \sum_{n \notin E_N, 1 \leq n \leq N} \frac{a^{2n}}{4\epsilon^2} \\ &\leq C^2 |E_N| + \frac{1}{4\epsilon^2(1-a^2)}. \end{aligned}$$

Because x is irrational, by Theorem 19 the sequence nx is uniformly distributed modulo 1. Therefore

$$\frac{|E_N|}{N} \rightarrow 2\epsilon, \quad N \rightarrow \infty,$$

and this implies

$$\limsup_{N \rightarrow \infty} \frac{S_N}{N} \leq 2\epsilon \cdot C^2.$$

Because this is true for each $0 < \epsilon < \frac{1}{2}$ it follows that $\lim_{N \rightarrow \infty} \frac{S_N}{N} = 0$, proving the claim. \square

Proposition 37. $R \leq \rho$.

Proof. We have found in (29) that $R \leq 2\rho$, i.e. $\rho \geq \frac{R}{2}$. Assume by contradiction that $R > \rho$; in particular $R > 0$, which implies $\rho \geq \frac{R}{2} > 0$. By Proposition 34, we have $F(z) = e^{f(z)}$ for $|z| < \rho$ and so $F(z) \neq 0$ for $z \in D(0, \rho)$.

Let $u_1\rho, \dots, u_s\rho$ be the distinct zeros of F on $|z| = \rho$, with respective multiplicities p_1, \dots, p_s ; if there is none, take $s = 0$, and use $\sum_{\emptyset} = 0$ and $\prod_{\emptyset} = 1$. Define

$$G(z) = F(z) \prod_{j=1}^s \left(1 - \frac{z}{u_j\rho}\right)^{-p_j}, \quad z \in D(0, R).$$

Because $G : D(0, R) \rightarrow \mathbb{C}$ is analytic and $G(z) \neq 0$ for $z \in \overline{D(0, \rho)}$, there is some T , $\rho < T \leq R$, such that $G(z) \neq 0$ for $z \in D(0, T)$ [130, p. 208, Theorem 10.18]. As $D(0, T)$ is simply connected, there is an analytic function $g : D(0, T) \rightarrow \mathbb{C}$ such that $G(z) = e^{g(z)}$ for $z \in D(0, T)$ [130, p. 274, Theorem 13.11]. Thus

$$F(z) = e^{g(z)} \prod_{j=1}^s \left(1 - \frac{z}{u_j\rho}\right)^{p_j}, \quad z \in D(0, T).$$

For $z \in D(0, \rho)$ we have $\left|\frac{z}{u_j\rho}\right| = \frac{|z|}{\rho} < 1$, and then by Lemma 32, $e^{L\left(\frac{z}{u_j\rho}\right)} = \left(1 - \frac{z}{u_j\rho}\right)^{-1}$. Therefore for $z \in D(0, \rho)$,

$$e^{f(z)} = F(z) = e^{g(z)} \prod_{j=1}^s e^{-p_j L\left(\frac{z}{u_j\rho}\right)} = \exp\left(g(z) - \sum_{j=1}^s p_j L\left(\frac{z}{u_j\rho}\right)\right),$$

i.e.

$$\exp\left(f(z) - g(z) - \sum_{j=1}^s p_j L\left(\frac{z}{u_j\rho}\right)\right) = 1, \quad z \in D(0, \rho).$$

But the image of a continuous function $D(0, \rho) \rightarrow 2\pi i\mathbb{Z}$ is connected, and because $2\pi i\mathbb{Z}$ has the discrete topology it follows that the image is a singleton, thus there is some $\nu \in \mathbb{Z}$ such that

$$f(z) - g(z) + \sum_{j=1}^s p_j L\left(\frac{z}{u_j\rho}\right) = 2\pi i\nu, \quad z \in D(0, \rho).$$

But $e^{g(0)} = G(0) = F(0) \cdot 1 = 1$, so $g(0) = 0$, and $f(0) = 0$, hence $\nu = 0$. Therefore

$$f(z) = g(z) - \sum_{j=1}^s p_j L\left(\frac{z}{u_j\rho}\right), \quad z \in D(0, \rho).$$

Now, for $z \in D(0, \rho)$,

$$\sum_{j=1}^s p_j L\left(\frac{z}{u_j\rho}\right) = \sum_{j=1}^s p_j \sum_{n=1}^{\infty} \frac{\left(\frac{z}{u_j\rho}\right)^n}{n} = \sum_{n=1}^{\infty} \left(\frac{1}{n} \sum_{j=1}^s p_j (u_j\rho)^{-n}\right) z^n.$$

Let $g_n = \frac{g^{(n)}(0)}{n!}$. For $z \in D(0, \rho)$,

$$\sum_{n=1}^{\infty} \gamma_n z^n = \sum_{n=1}^{\infty} g_n z^n - \sum_{n=1}^{\infty} \left(\frac{1}{n} \sum_{j=1}^s p_j (u_j \rho)^{-n} \right) z^n,$$

so for $n \geq 1$,

$$\frac{1}{n(1 - q_0^n)} = \gamma_n = g_n - \frac{1}{n} \sum_{j=1}^s p_j (u_j \rho)^{-n}.$$

Then

$$\frac{\rho^n}{1 - q_0^n} = n \rho^n g_n - \sum_{j=1}^s p_j u_j^{-n}. \quad (30)$$

Cauchy's integral formula [144, p. 82, Theorem 2.41] tells us that for $0 < V < T$, $C(t) = V e^{it}$, $0 \leq t \leq 2\pi$,

$$g_n = \frac{g^{(n)}(0)}{n!} = \frac{1}{2\pi i} \int_C \frac{g(w)}{w^{n+1}} dw,$$

whence, as the length of C is $2\pi V$,

$$|g_n| \leq \frac{1}{V^n} \max_{|w|=V} |g(w)|.$$

Fix $\rho < V < T$, with which

$$|n \rho^n g_n| \leq n \left(\frac{\rho}{V} \right)^n \max_{|w|=V} |g(w)|,$$

and for $\frac{\rho}{V} < \delta < 1$ we have

$$|n \rho^n g_n| = O(\delta^n).$$

Using this and

$$\left| \sum_{j=1}^s p_j u_j^{-n} \right| \leq \sum_{j=1}^s p_j = O(1),$$

(30) yields

$$\frac{\rho^n}{1 - q_0^n} = O(1).$$

As

$$\frac{\rho^n}{1 - q_0^n} = \frac{\rho^n}{1 - e^{2\pi i n x}} = \frac{\rho^n}{-2ie^{\pi i n x} \sin \pi n x} = \frac{ie^{-\pi i n x} \rho^n}{2 \sin \pi n x},$$

we get

$$\frac{\rho^{2n}}{\sin^2 \pi n x} = O(1).$$

For any M , there is some n_M such that $\sin^2 \pi n_M x \geq M$, and thus the above estimate is contradicted if $\rho = 1$; hence $0 < \rho < 1$. (We emphasize that $\rho < 1$ is deduced from the assumption $R > \rho$, which we are showing to imply a contradiction.) By Lemma 36 we then get that

$$\sum_{n=1}^N \frac{\rho^{2n}}{\sin^2 n\pi x} = o(N). \quad (31)$$

Now, multiplying each side of (30) by its complex conjugate and using that $|n\rho^n g_n| = O(\delta^n)$ and that $|\sum_{j=1}^s p_j u_j^{-n}| = O(1)$,

$$\frac{\rho^{2n}}{4 \sin^2 \pi n x} = \left(\sum_{j=1}^s p_j u_j^{-n} \right) \left(\sum_{k=1}^s p_k u_k^n \right) + O(\delta^n),$$

i.e.

$$\frac{\rho^{2n}}{4 \sin^2 \pi n x} = \sum_{j=1}^s p_j^2 + \sum_{j \neq k} p_j p_k (u_j^{-1} u_k)^n + O(\delta^n). \quad (32)$$

Let $E = \{(j, k) : 1 \leq j, k \leq s, j \neq k\}$ and let $P = \sum_{j=1}^s p_j^2 > 0$. Then summing (32) for $n = 1, \dots, N$, using $\sum_{n=1}^N \delta^n = \delta \cdot \frac{1-\delta^N}{1-\delta} = O(1)$,

$$\sum_{n=1}^N \frac{\rho^{2n}}{4 \sin^2 \pi n x} = NP + \sum_{(j,k) \in E} p_j p_k \left(\sum_{n=1}^N (u_j^{-1} u_k)^n \right) + O(1).$$

Because $u_j \neq u_k$ for $(j, k) \in E$, we have according to Lemma 35 that

$$\left| \sum_{n=1}^N (u_j^{-1} u_k)^n \right| \leq \frac{2}{|1 - u_j^{-1} u_k|} = O(1).$$

Thus

$$\sum_{n=1}^N \frac{\rho^{2n}}{\sin^2 \pi n x} = 4NP + O(1),$$

and because $P > 0$ this contradicts (31). Therefore, it is false that $R > \rho$, which means that $R \leq \rho$, proving the claim. \square

Theorem 38. *Let $x \in \Omega$ and let ρ_1 and R_1 respectively be the radii of convergence of the power series*

$$\sum_{n=1}^{\infty} \frac{z^n}{\sin n\pi x}, \quad \sum_{n=1}^{\infty} \frac{z^n}{\sin \pi x \cdot \sin 2\pi x \cdots \sin n\pi x}.$$

Then $R_1 = \frac{\rho_1}{2}$.

Proof.

$$|1 - q_0^n| = |1 - e^{2\pi i n x}| = 2|\sin \pi n x|.$$

By the Cauchy-Hadamard formula,

$$R_1 = \liminf_{n \rightarrow \infty} |\sin \pi x \cdots \sin n \pi x|^{1/n} = \frac{1}{2} \cdot \liminf_{n \rightarrow \infty} |(1 - q_0) \cdots (1 - q_0^n)|^{1/n} = \frac{R}{2}$$

and

$$\rho_1 = \liminf_{n \rightarrow \infty} |\sin n \pi x|^{1/n} = \rho.$$

Theorem 30 says $R = \rho$, hence $R_1 = \frac{R}{2} = \frac{\rho}{2} = \frac{\rho_1}{2}$. \square

By Lemma 23 and Theorem 24, for almost all x , the power series $\sum_{n=1}^{\infty} \frac{z^n}{\sin n \pi x}$ has radius of convergence 1. Then using Theorem 38, for almost all x the power series $\sum_{n=1}^{\infty} \frac{z^n}{\sin \pi x \cdot \sin 2\pi x \cdots \sin n \pi x}$ has radius of convergence $\frac{1}{2}$, and thus for almost all $x \in \Omega$,

$$\liminf_{n \rightarrow \infty} \left(\prod_{k=1}^n |\sin k \pi x| \right)^{1/n} = \frac{1}{2}.$$

Hardy and Littlewood give a separate argument [60, p. 88, Eq. 4.3] proving that for almost all $x \in \Omega$,

$$\limsup_{n \rightarrow \infty} \left(\prod_{k=1}^n |\sin k \pi x| \right)^{1/n} = \frac{1}{2},$$

and combining the formulas for the limit inferior and limit superior yields Theorem 29.

Lubinsky [98] proves more results about products of the form $\prod_{k=1}^n (1 - e^{2\pi i k \theta})$. For example, Lubinsky [98, p. 219, Theorem 1.1] proves that for all $\epsilon > 0$, for almost all $\theta \in \Omega$ we have

$$\left| \log \left| \prod_{k=1}^n (1 - e^{2\pi i k \theta}) \right| \right| = O((\log n)(\log \log n)^{1+\epsilon}).$$

The author [14, p. 532, Theorem 2] gives asymptotic expressions for the $L^p[0, 1]$ norm of $\prod_{k=1}^n (1 - e^{2\pi i k \theta})$ as $n \rightarrow \infty$, for $1 \leq p \leq \infty$.

For $\omega \in \mathbb{R}$ let $P_n(\omega) = \prod_{r=1}^n |2 \sin \pi r \omega|$. Let F_n be the n th Fibonacci number and let $g = \frac{-1 + \sqrt{5}}{2}$. Verschueren and Mestel [149, p. 204, Theorem 2.2] prove that there is some $c = 2.407 \dots$ such that

$$P_{F_n}(g) \rightarrow c, \quad n \rightarrow \infty$$

and

$$\frac{P_{F_{n-1}}(g)}{F_n} \rightarrow \frac{c}{2\pi\sqrt{5}}, \quad n \rightarrow \infty,$$

and that there are $C_1 \leq 0$ and $C_2 \geq 1$ such that for all n ,

$$n^{C_1} \leq P_n(g) \leq n^{C_2}.$$

Let X be a measure space with probability measure λ . Following [138, p. 21, Definition 3.6], we say that a measure preserving map $T : X \rightarrow X$ is **r -fold mixing** if for all $g, f_1, \dots, f_r \in L^{r+1}(X)$ we have

$$\begin{aligned} & \lim_{m_1 \rightarrow \infty, \dots, m_r \rightarrow \infty} \int_X g(t) \cdot \prod_{k=1}^r f_k \left(T^{\sum_{j=1}^k m_j}(t) \right) d\lambda(t) \\ &= \left(\int_X g(t) d\lambda(t) \right) \prod_{k=1}^r \left(\int_X f_k(t) d\lambda(t) \right). \end{aligned} \tag{33}$$

If for each r the map T is r -fold mixing, we say that T is **mixing of all orders**.

Let $q \geq 2$ be an integer, and define $T_q : [0, 1] \rightarrow [0, 1]$ by $T_q(t) = R(qt)$. We assert that T_q is mixing of all orders. This can be proved by first showing that the dynamical system $([0, 1], \mu, T_q)$ is isomorphic to a Bernoulli shift (cf. [44, p. 17, Example 2.8]). This implies that if the Bernoulli shift is r -fold mixing then T_q is r -fold mixing. One then shows that a Bernoulli shift is mixing of all orders [44, p. 53, Exercise 2.7.9]. Using that T_q is mixing of all orders gets us the following result.

Theorem 39. *Let $q \geq 2$ be an integer. For each $n \geq 1$ we have*

$$\lim_{m \rightarrow \infty} \int_0^1 |\sin(2\pi t)| \cdot \prod_{k=1}^n |\sin(2\pi q^{km} t)| dt = \left(\frac{2}{\pi} \right)^{n+1}.$$

Proof. Define $g(t) = f_1(t) = \dots = f_n(t) = |\sin(2\pi t)|$. For any nonzero integer N we have

$$\int_0^1 |\sin(2\pi Nt)| dt = \frac{2}{\pi},$$

and it follows from (33), using $m_1 = m, \dots, m_n = m$, that

$$\lim_{m \rightarrow \infty} \int_0^1 |\sin(2\pi t)| \cdot \prod_{k=1}^n |\sin(2\pi q^{km} t)| dt = \left(\frac{2}{\pi} \right)^{n+1}.$$

□

See Sinai [137].

Write $S_k(\alpha) = \sum_{j=1}^k X_j(\alpha)$, where $X_j(\alpha) = \log |2 - 2\cos(2\pi j\alpha)|$ and $\alpha = \frac{-1+\sqrt{5}}{2}$. Knill and Tangerman [79] talk about motivations from KAM theory for caring about these sums. See Lagarias [86] and Ghys [54] for more on small divisors in Hamiltonian dynamics, and Carleson and Gamelin [26, p. 48, Theorem 7.2] and Yoccoz [159] for Arnold's theorem on analytic circle diffeomorphisms.

Marmi and Sauzin [99]

12 Conclusions

Kac and Salem [73] prove the following. Let c_k be a sequence of nonnegative real numbers for which $\sum_{k=1}^{\infty} c_k < \infty$. If the series

$$\sum_{k=1}^{\infty} c_k \frac{1}{|\sin kx|}$$

converges in a set of positive measure, then

$$\sum_{k=1}^{\infty} c_k \log \left(\frac{1}{c_k} \right) < \infty,$$

and if this condition is satisfied then $\sum_{k=1}^{\infty} c_k \frac{1}{|\sin kx|}$ converges for almost all x .

Muromskii [105, p. 54, Theorem 1] proves that if c_k is a sequence of nonnegative real numbers, if $\alpha > 1$, and if there is a set of positive measure on which the series

$$\sum_{k=1}^{\infty} c_k \frac{1}{|\sin kx|^\alpha}$$

converges, then for any $\delta > 0$, the series

$$\sum_{k=1}^{\infty} c_k^{\frac{4}{\alpha+3}+\delta}$$

converges.

Let X be a random variable that is uniformly distributed on $[0, 1]$. Kesten [77, p. 111, Theorem 1] proves that if $\sum_{k=0}^{\infty} |c_k| < \infty$, then the series

$$\sum_{k=0}^{\infty} \frac{c_k}{\sin(2\pi 2^k X)}$$

converges with probability 1. Stated using measure theory, the conclusion is that for almost all $x \in \Omega$, the series

$$\sum_{k=0}^{\infty} \frac{c_k}{\sin(2\pi 2^k x)}$$

converges. Kesten [77, p. 114, Theorem 3] also proves if $a_n \in \mathbb{R}$ and $a_n \rightarrow \infty$, then

$$\frac{1}{a_n} \sum_{k=0}^{n-1} \frac{1}{\sin(2\pi 2^k X)} \rightarrow 0$$

in probability. Stated using measure theory, the conclusion is that for each $\epsilon > 0$,

$$\lim_{n \rightarrow \infty} \mu \left\{ x \in \Omega : \left| \frac{1}{a_n} \sum_{k=0}^{n-1} \frac{1}{\sin(2\pi 2^k x)} \right| \geq \epsilon \right\} = 0.$$

For $\mathbb{T}^d = \mathbb{R}^d/\mathbb{Z}^d = (\mathbb{R}/\mathbb{Z})^d$, let σ be Haar measure on \mathbb{T} and let $\sigma_d = \bigotimes_{1 \leq j \leq d} \sigma$ be Haar measure on \mathbb{T}^d , with $\sigma(\mathbb{T}) = 1$. A sequence $t(n) \in \mathbb{T}^d$, $n \geq 1$, is said to be **uniformly distributed** if for any arcs I_1, \dots, I_d in \mathbb{T} , with $I = \prod_{j=1}^d I_j$,

$$\lim_{N \rightarrow \infty} \frac{|\{t(n) \in I : 1 \leq n \leq N\}|}{N} = \sigma_d(I).$$

Kronecker's approximation theorem [145, p. 108, Theorem 6.3] states that if $\alpha_1, \dots, \alpha_d \in \mathbb{R}$ and $\{1, \alpha_1, \dots, \alpha_d\}$ is linearly independent over \mathbb{Q} , then the sequence $(n\alpha_1 + \mathbb{Z}, \dots, n\alpha_d + \mathbb{Z})$, $n \geq 1$, is uniformly distributed in \mathbb{T}^d . Meyer [100] is a thorough presentation of multidimensional Diophantine approximation and Diophantine approximation with locally compact abelian groups, and harmonic analysis involving sets satisfying various Diophantine properties.

See Rudin [128].

See Abraham and Marsden [1, p. 395, Proposition 5.2.23] and [1, pp. 818–820, Proposition 5.2.23].

Measure theoretic results in Diophantine approximation are presented in Khinchin [78], Einsiedler and Ward [44, Chapter 3], Rockett and Szűs [127, Chapters V and VI], Billingsley [19, pp. 13–15, 319–326], Kac [74, Chapter 5], Bugeaud [25], and Kesseböhmer, Munday and Stratmann [76]. The significance of continued fraction expansions of irrational numbers in the early history of axiomatic probability theory is described by Barone and Novikoff [8], Durand and Mazliak [43], and von Plato [152]. Veech [148] presents material on Diophantine approximation in the setting of topological dynamics.

We have been interested in results about almost all $x \in \Omega$, using Lebesgue measure μ on $[0, 1]$. If $E \subset \Omega$ has $\mu(E) = 0$, one can ask what the Hausdorff dimension $\dim_H E$ of the set E is. Let \mathcal{B}_K be the set of those $x \in \Omega$ such that $a_n(x) \leq K$ for all $n \geq 1$, and let $\mathcal{B} = \bigcup_{K \geq 1} \mathcal{B}_K$, the set of those $x \in \Omega$ with bounded partial quotients. We have already stated that $\mu(\mathcal{B}) = 0$ [78, p. 60, Theorem 29], and Jarník [39, Theorem 4.3] proves that the Hausdorff dimension of \mathcal{B} is in fact 1; cf. Falconer [47, p. 155, Theorem 10.3] and Wolff [158, p. 67, Chapter 9] on Hausdorff dimension. Hensley [64] proves

$$\dim_H \mathcal{B}_K = 1 - \frac{6}{\pi^2} K^{-1} - \frac{72}{\pi^4} K^{-2} \log K + O(K^{-2}), \quad K \rightarrow \infty.$$

Dodson and Kristensen [39] give a survey of results on the Hausdorff dimension of various sets that appear in Diophantine approximation.

For a decreasing positive function ψ , the set

$$W(\psi) = \{x \in [0, 1] : \|qx\| < q\psi(q) \text{ for infinitely many } q \in \mathbb{N}\}$$

can be written as the limsup of a sequence of sets,

$$W(\psi) = \limsup_{n \rightarrow \infty} W(\psi, n) = \bigcap_{N=1}^{\infty} \bigcup_{n=N}^{\infty} W(\psi, n),$$

where

$$W(\psi, n) = \bigcup_{2^{n-1} < q \leq 2^n} \bigcup_{0 \leq p \leq q} \left(\frac{p}{q} - \psi(q), \frac{p}{q} + \psi(q) \right) \cap [0, 1].$$

One can exploit nice properties of limsup sets, such as the Borel-Cantelli lemma and invariance under ergodic transformations, to prove fundamental results in Diophantine approximation. Beresnevich, Dickinson and Velani [16] use this motivation of Diophantine approximation to build a framework for a natural class of limsup sets on compact metric spaces. Their general results readily imply the divergent case of Khinchin's theorem: $\mu(W(\psi)) = 0$ if $\sum q\psi(q) < \infty$ (Lemma 7) and $\mu(W(\psi)) = 1$ if $\sum q\psi(q) = \infty$. Their framework also establishes the divergent case of Jarník's theorem: the f -Hausdorff dimension of $W(\psi)$ is 0 if $\sum qf(\psi(q)) < \infty$ and is infinity if $\sum qf(\psi(q)) = \infty$, where f is a **dimension function** such that $r^{-1}f(r) \rightarrow \infty$ as $r \rightarrow 0$, and $r^{-1}f(r)$ is decreasing.

As well, rather than making statements about subsets of Ω of measure 1, we can talk about sets whose complements are meager. (Measure theoretically, the notion of a negligible set is made precise as a set of measure 0, and topologically the notion of a negligible set is made precise as a meager set.) Some results of this type are proved in Oxtoby [111, Chapter 2].

Let p be prime, let $N_p = \{0, \dots, p-1\}$, and let $\mathbb{Q}_p \subset \prod_{\mathbb{Z}} N_p$ be the p -adic numbers. For $x \in \mathbb{Q}_p$ let

$$v_p(x) = \inf\{k \in \mathbb{Z} : x(k) \neq 0\}, \quad |x|_p = p^{-v_p(x)},$$

and let $\mathbb{Z}_p = \{x \in \mathbb{Q}_p : v_p(x) \geq 0\}$, the p -adic integers. Let μ_p be the Haar measure on the additive locally compact abelian group \mathbb{Q}_p with $\mu_p(\mathbb{Z}_p) = 1$. We call $\lambda \in \mathbb{Z}_p$ a **p -adic Liouville number** if $\nu(\lambda) = \liminf_{n \rightarrow \infty} |n - \lambda|_p^{1/n} = 0$, and let \mathcal{L}_p be the set of p -adic Liouville numbers. One checks that if $\lambda \in \mathbb{Z}_{\geq 0}$ then $\nu(\lambda) = 1$ [131, p. 201, Exercise 66.A]. It can be proved that \mathcal{L}_p is a dense G_δ set in \mathbb{Z}_p [131, p. 204, Theorem 67.3] and that $\mu_p(\mathcal{L}_p) = 0$ [131, p. 205, Theorem 67.4]. One reason for caring about p -adic Liouville numbers is that if $x \in \mathbb{Z}_p$ is algebraic over \mathbb{Q} then $\nu(x) = 1$, and hence a p -adic Liouville number is transcendental over \mathbb{Q} [131, p. 203, Theorem 67.2].

Unlike in estimating exponential sums, the sums that we have been estimating in this paper do not have cancellation. Instead we have estimated them by showing that the terms are only occasionally large. For $A_n(t) = \sum_{k=1}^n \frac{\sin kt}{k}$, it can be proved [118, p. 74, no. 25] that

$$|A_n(t)| < \int_0^\pi \frac{\sin \theta}{\theta} d\theta = 1.8519\dots$$

On the other hand, let M_n be the maximum of $\sum_{k=1}^n \frac{|\sin kt|}{k}$. It can be proved [118, p. 77, no. 38] that

$$M_n = \frac{2}{\pi} \log n + O(1).$$

Walfisz [153] presents results of his, of Oppenheim, and of Chowla on sums

$$\sum_{j \leq n} g(j) e^{2\pi i j x}$$

for $g(j) = r_k(j)$, the number of ways to write j as a sum of k squares, and for $g(j) = d(j)$, the number of positive divisors of j . One of the results of Chowla is that if $x \in \Omega$ has bounded partial quotients, then

$$\sum_{j=1}^n d(j)e^{2\pi i j x} = O\left(n^{\frac{1}{2}} \log n\right).$$

One of the results Walfisz proves is that if $\epsilon > 0$, then for almost all $x \in \Omega$,

$$\sum_{j=1}^n d(j)e^{2\pi i j x} = O\left(n^{\frac{1}{2}}(\log n)^{2+\epsilon}\right).$$

Wilton [157] proves some similar results. For example, Wilton proves that for any $x \in \Omega$,

$$\sum_{j=1}^n \frac{d(j)}{j} \cos 2\pi j x = o((\log n)^2),$$

See Jutila's book on exponential sums [72].

Let $f(t) = P_1(t)$ for $t \notin \mathbb{Z}$ and $\{t\} = 0$ for $t \in \mathbb{Z}$, where P_1 is a periodic Bernoulli function. Namely, for all $t \in \mathbb{R}$,

$$f(t) = -\frac{1}{\pi} \sum_{m=1}^{\infty} \frac{\sin 2\pi m t}{m}.$$

Define $S_N(\theta) = \sum_{n=1}^N \frac{\mu(n)}{n} f(n\theta)$, where μ is the **Möbius function**. Davenport [33, p. 11, Theorem 2] proves that there is some C such that for all N and for all θ ,

$$\left| \sum_{n=1}^N \frac{\mu(n)}{n} f(n\theta) \right| \leq C.$$

Davenport [33, p. 13, Theorem 4] also proves that for almost all θ ,

$$\sum_{n=1}^N \frac{\mu(n)}{n} f(n\theta) \rightarrow -\frac{1}{\pi} \sin 2\pi\theta.$$

See Jaffard [70].

It would be a useful project to give an organized presentation of Hardy and Littlewood's results on Diophantine approximation. Their papers in this area are all included in Hardy's collected works [58]. Hardy and Littlewood proved many pleasant results on various sums and series with coefficients related to $\sin(n\pi x)$ and $R(nx)$. It would be desirable to streamline and systematically prove these results, to let a modern reader to be able to understand them without having to read the whole series of papers to figure out what results are being tacitly used from earlier work or assumed as general knowledge. There is only a bare summary of Hardy and Littlewood's work in the commentary in Hardy's collected papers. Hardy's work on Diophantine approximation is briefly summarized by Mordell [104]. See also lecture V of Hardy's lectures on Ramanujan [57].

Acknowledgments

The author thanks Hervé Queffélec (Université de Lille) and Martine Queffélec (Université de Lille) for a long correspondence on Hardy and Littlewood’s “curious power-series” [60].

The author thanks Jeremy Voltz (University of Toronto) for discussions on Beresnevich, Dickinson and Velani’s monograph [16] on limsup sets.

References

- [1] Ralph Abraham and Jerrold E. Marsden. *Foundations of Mechanics*. AMS Chelsea Publishing, Providence, Rhode Island, second edition, 2008.
- [2] Charalambos D. Aliprantis and Kim C. Border. *Infinite Dimensional Analysis: A Hitchhiker’s Guide*. Springer, third edition, 2006.
- [3] George E. Andrews. *The Theory of Partitions*, volume 2 of *Encyclopedia of Mathematics and Its Applications*. Addison-Wesley, 1976.
- [4] Tom M. Apostol. *Modular forms and Dirichlet series in number theory*, volume 41 of *Graduate Texts in Mathematics*. Springer, second edition, 1990.
- [5] L. C. Arboleda. Les recherches de M. Fréchet, P. Alexandrov, W. Sierpinski et K. Kuratowski sur la théorie des types de dimensions et les débuts de la topologie générale. *Arch. Hist. Exact Sci.*, 24(4):339–388, 1981.
- [6] Paul Bachmann. Analytische Zahlentheorie. In Wilhelm Franz Meyer, editor, *Encyklopädie der mathematischen Wissenschaften mit Einschluss ihrer Anwendungen, Band I, 1. Teil*, pages 636–674. B. G. Teubner, Leipzig, 1898–1904.
- [7] Paul Bachmann. Über Gauss’ zahlentheoretische Arbeiten. In Gesellschaft der Wissenschaften zu Göttingen, editor, *Carl Friedrich Gauss Werke, Band 10, Abteilung 2*, pages 1–74. Julius Springer, Berlin, 1922–1933.
- [8] Jack Barone and Albert Novikoff. A history of the axiomatic formulation of probability from Borel to Kolmogorov: Part I. *Arch. Hist. Exact Sci.*, 18(2):123–190, 1978.
- [9] Christoph Baxa. Calculation of improper integrals using uniformly distributed sequences. *Acta Arith.*, 119(4):373–406, 2005.
- [10] József Beck. From probabilistic Diophantine approximation to quadratic fields. In Peter Hellekalek and Gerhard Larcher, editors, *Random and Quasi-Random Point Sets*, volume 138 of *Lecture Notes in Statistics*, pages 1–48. Springer, 1998.

- [11] József Beck. *Probabilistic Diophantine Approximation: Randomness in Lattice Point Counting*. Springer, 2014.
- [12] József Beck and William W. L. Chen. *Irregularities of distribution*. Number 89 in Cambridge Tracts in Mathematics. Cambridge University Press, 1987.
- [13] Heinrich Behnke. Zur Theorie der diophantischen Approximationen. *Abhandlungen aus dem mathematischen Seminar der Hamburgischen Universität*, 3:261–318, 1924.
- [14] Jordan Bell. Estimates for the norms of products of sines and cosines. *J. Math. Anal. Appl.*, 405(2):530–545, 2013.
- [15] John J. Benedetto and Wojciech Czaja. *Integration and Modern Analysis*. Birkhäuser Advanced Texts. Birkhäuser, 2009.
- [16] Victor Beresnevich, Detta Dickinson, and Sanju Velani. *Measure theoretic laws for lim sup sets*. Number 846 in Memoirs of the American Mathematical Society. American Mathematical Society, Providence, RI, 2006.
- [17] Bruce C. Berndt. Dedekind sums and a paper of G. H. Hardy. *J. London Math. Soc. (2)*, 13(1):129–137, 1976.
- [18] Valérie Berthé. Autour du système de numération d’Ostrowski. *Bull. Belg. Math. Soc. Simon Stevin*, 8(2):209–239, 2001.
- [19] Patrick Billingsley. *Probability and Measure*. John Wiley & Sons, third edition, 1995.
- [20] Harald Bohr and Harald Cramér. Die neue Entwicklung der analytischen Zahlentheorie. In H. Burkhardt, W. Wirtinger, R. Fricke, and E. Hilb, editors, *Encyklopädie der mathematischen Wissenschaften mit Einschluss ihrer Anwendungen, Band II, 3. Teil, 2. Hälfte*, pages 722–849. B. G. Teubner, Leipzig, 1923–1927.
- [21] Francis Bonahon. *Low-Dimensional Geometry: From Euclidean Surfaces to Hyperbolic Knots*, volume 49 of *Student Mathematical Library*. American Mathematical Society, Providence, RI, 2009.
- [22] Jonathan Breuer and Barry Simon. Natural boundaries and spectral theory. *Adv. Math.*, 226(6):4902–4920, 2011.
- [23] Claude Brezinski. *History of Continued Fractions and Padé Approximants*. Springer, 1991.
- [24] Tom C. Brown and Peter Jau-Shyong Shiue. Sums of fractional parts of integer multiples of an irrational. *J. Number Theory*, 50(2):181–192, 1995.

- [25] Yann Bugeaud. *Distribution modulo one and Diophantine approximation*. Number 193 in Cambridge Tracts in Mathematics. Cambridge University Press, 2012.
- [26] Lennart Carleson and Theodore W. Gamelin. *Complex dynamics*. Universitext. Springer, 1993.
- [27] J. W. S. Cassels. *An Introduction to Diophantine Approximation*. Number 45 in Cambridge Tracts in Mathematics and Mathematical Physics. Cambridge University Press, 1957.
- [28] Ernesto Cesàro. Question 1151. *L'Intermédiaire des mathématiciens*, 4:220–221, 1897.
- [29] K. Chandrasekharan. Exponential sums in the development of number theory. In I. M. Vinogradov, A. A. Karacuba, K. I. Oskolkov, and A. N. Paršin, editors, *Proceedings of the International Conference on Number Theory (Moscow, September 14–18, 1971)*, volume 132 of *Trudy Mat. Inst. Steklov.*, pages 7–26. Nauka, 1973.
- [30] I. P. Cornfeld, S. V. Fomin, and Ya. G. Sinai. *Ergodic Theory*, volume 245 of *Grundlehren der mathematischen Wissenschaften*. Springer, 1982. Translated by A. B. Sossinskii.
- [31] Rosanna Cretney. The origins of Euler’s early work on continued fractions. *Historia Math.*, 41(2):139–156, 2014.
- [32] Joseph Warren Dauben. *Georg Cantor: His Mathematics and Philosophy of the Infinite*. Princeton University Press, 1979.
- [33] Harold Davenport. On some infinite series involving arithmetical functions. *Q. J. Math.*, 8:8–13, 1937. Collected Works, vol. IV, pp. 1781–1786.
- [34] Charles-Jean de la Vallée-Poussin. Question 1490. *L'Intermédiaire des mathématiciens*, 6:213–214, 1899.
- [35] Leonard Eugene Dickson. *History of the theory of numbers, volume I: Divisibility and primality*. Carnegie Institution of Washington, Washington, 1919.
- [36] Leonard Eugene Dickson. *History of the theory of numbers, volume III: Quadratic and higher forms*. Carnegie Institution of Washington, Washington, 1923. With a chapter on the class number by G. H. Cresse.
- [37] P. G. L. Dirichlet. Über die Bestimmung der mittleren Werthe in der Zahlentheorie. *Abhandlungen der Königl. Akademie der Wissenschaften zu Berlin*, pages 69–83, 1849. Werke, Band II, pp. 49–66.

- [38] P. G. L. Dirichlet. *Lectures on Number Theory*, volume 16 of *History of Mathematics*. American Mathematical Society, Providence, RI, 1999. Supplements by R. Dedekind, translated from the German by John Stillwell.
- [39] M. Maurice Dodson and Simon Kristensen. Hausdorff dimension and Diophantine approximation. In Michel L. Lapidus and Machiel van Frankenhuysen, editors, *Fractal Geometry and Applications: A Jubilee of Benoît Mandelbrot, Part 1*, volume 72 of *Proceedings of Symposia in Pure Mathematics*, pages 305–347. American Mathematical Society, Providence, RI, 2004.
- [40] K. A. Driver, D. S. Lubinsky, G. Petruska, and P. Sarnak. Irregular distribution of $\{n\beta\}$, $n = 1, 2, 3, \dots$, quadrature of singular integrands, and curious basic hypergeometric series. *Indag. Math. (N.S.)*, 2(4):469–481, 1991.
- [41] Michael Drmota and Robert F. Tichy. *Sequences, discrepancies and applications*, volume 1651 of *Lecture Notes in Mathematics*. Springer, 1997.
- [42] William Duke and Özlem Imamoğlu. Lattice points in cones and Dirichlet series. *Int. Math. Res. Not.*, (53):2823–2836, 2004.
- [43] Antonin Durand and Laurent Mazliak. Revisiting the sources of Borel’s interest in probability: continued fractions, social involvement, Volterra’s prolusione. *Centaurus*, 53(4):306–332, 2011.
- [44] Manfred Einsiedler and Thomas Ward. *Ergodic Theory: with a view towards Number Theory*, volume 259 of *Graduate Texts in Mathematics*. Springer, 2010.
- [45] Paul Erdős. Some remarks on Diophantine approximations. *J. Indian Math. Soc. (N.S.)*, 12:67–74, 1948.
- [46] Leonhard Euler. De summis serierum numeros Bernoullianos involventium. *Novi Commentarii Academiae Scientiarum Imperialis Petropolitanae*, 14:129–167, 1770. E393, Opera omnia I.15, pp. 91–130.
- [47] Kenneth Falconer. *Fractal geometry: mathematical foundations and applications*. John Wiley & Sons, second edition, 2003.
- [48] N. I. Feldman and Yu. V. Nesterenko. *Number theory IV: Transcendental numbers*, volume 44 of *Encyclopaedia of Mathematical Sciences*. Springer, 1998. Translated from the Russian by Neal Koblitz.
- [49] José Ferreirós. *Labyrinth of Thought: A History of Set Theory and Its Role in Modern Mathematics*, volume 23 of *Science Networks. Historical Studies*. Birkhäuser, second revised edition, 2007.

- [50] David Fowler. Newton, Cotes, and $\sqrt{\sqrt{2}}$: a footnote to Newton's theory of the resistance of fluids. In P. M. Harman and Alan E. Shapiro, editors, *The Investigation of Difficult Things. Essays on Newton and the History of the Exact Sciences in Honour of D. T. Whiteside*, pages 355–368. Cambridge University Press, 1992.
- [51] David Fowler. *The mathematics of Plato's Academy: a new reconstruction*. Clarendon Press, Oxford, second edition, 1999.
- [52] Jérôme Franel. Question 1260. *L'Intermédiaire des mathématiciens*, 5:77, 1898.
- [53] Jérôme Franel. Question 1547. *L'Intermédiaire des mathématiciens*, 6:149–150, 1899.
- [54] Étienne Ghys. Resonances and small divisors. In Éric Charpentier, Annick Lesne, and Nikolai K. Nikolski, editors, *Kolmogorov's Heritage in Mathematics*, pages 187–213. Springer, 2007.
- [55] Ronald L. Graham, Donald E. Knuth, and Oren Patashnik. *Concrete Mathematics: A Foundation for Computer Science*. Addison-Wesley, second edition, 1994.
- [56] Seymour Haber and Charles F. Osgood. On the sum $\sum \langle n\alpha \rangle^{-t}$ and numerical integration. *Pacific J. Math.*, 31(2):383–394, 1969.
- [57] G. H. Hardy. *Ramanujan: twelve lectures on subjects suggested by his life and work*. Cambridge University Press, 1940.
- [58] G. H. Hardy. *Collected Papers of G. H. Hardy*, volume I. Oxford University Press, 1966.
- [59] G. H. Hardy and J. E. Littlewood. Some problems of Diophantine approximation: The lattice-points of a right-angled triangle. *Proc. London Math. Soc., Ser. 2*, 20:15–36, 1922. Collected Papers, vol. I, pp. 136–158.
- [60] G. H. Hardy and J. E. Littlewood. Notes on the theory of series (XXIV): a curious power-series. *Proc. Cambridge Philos. Soc.*, 42(2):85–90, 1946. Collected Papers, vol. I, pp. 253–259.
- [61] G. H. Hardy and E. M. Wright. *An introduction to the theory of numbers*. Clarendon Press, Oxford, fifth edition, 1979.
- [62] Masayoshi Hata. *Problems and Solutions in Real Analysis*. World Scientific, 2007.
- [63] Thomas Hawkins. *Lebesgue's theory of integration: its origins and development*. AMS Chelsea Publishing, Providence, RI, second edition, 2001.
- [64] Doug Hensley. Continued fraction Cantor sets, Hausdorff dimension, and functional analysis. *J. Number Theory*, 40:336–358, 1992.

- [65] Ch. Hermite. Sur quelques conséquences arithmétiques des formules de la théorie des fonctions elliptiques. *Acta Math.*, 5(1):297–330, 1884. Œuvres, tome IV, pp. 138–168.
- [66] Edmund Hlawka and Christa Binder. Über die Entwicklung der Theorie der Gleichverteilung in den Jahren 1909 bis 1916. *Arch. Hist. Exact Sci.*, 36(3):197–249, 1986.
- [67] Edmund Hlawka, Johannes Schoißengeier, and Rudolf Taschner. *Geometric and analytic number theory*. Universitext. Springer, 1991. Translated from the German by Charles Thomas.
- [68] Loo-Keng Hua. Die Abschätzung von Exponentialsummen und ihre Anwendung in der Zahlentheorie. In M. Deuring, H. Hasse, and E. Sperner, editors, *Enzyklopädie der mathematischen Wissenschaften mit Einschluss ihrer Anwendungen, Band II, 2. Teil, Heft 13, Teil I*, pages 1–123. B. G. Teubner, Leipzig, 1959.
- [69] Marius Iosifescu and Cor Kraaikamp. *Metrical theory of continued fractions*, volume 547 of *Mathematics and Its Applications*. Kluwer Academic Publishers, 2002.
- [70] Stéphane Jaffard. On Davenport expansions. In Michel L. Lapidus and Machiel van Frankenhuysen, editors, *Fractal Geometry and Applications: A Jubilee of Benoît Mandelbrot, Part 1*, volume 72 of *Proceedings of Symposia in Pure Mathematics*, pages 273–303. American Mathematical Society, Providence, RI, 2004.
- [71] Dale M. Johnson. The problem of the invariance of dimension in the growth of modern topology, part II. *Arch. Hist. Exact Sci.*, 25(2–3):85–266, 1981.
- [72] M. Jutila. *Lectures on A Method in the Theory of Exponential Sums*, volume 80 of *Tata Institute of Fundamental Research Lectures on Mathematics*. Springer, 1987.
- [73] M. Kac and R. Salem. On a series of cosecants. *Nederl. Akad. Wetensch. Proc. Ser. A*, 60:265–267, 1957.
- [74] Mark Kac. *Statistical independence in probability, analysis and number theory*. Number 12 in Carus Mathematical Monographs. Mathematical Association of America, 1959.
- [75] Alexander S. Kechris. *Classical descriptive set theory*, volume 156 of *Graduate Texts in Mathematics*. Springer, 1995.
- [76] Marc Kesseböhmer, Sara Munday, and Bernd O. Stratmann. *Infinite Ergodic Theory of Numbers*. De Gruyter, 2016.

- [77] Harry Kesten. On a series of cosecants. II. *Nederl. Akad. Wetensch. Proc. Ser. A*, 62:110–119, 1959.
- [78] A. Ya. Khinchin. *Continued fractions*. Dover Publications, third edition, 1997. Translated from the Russian.
- [79] Oliver Knill and Folkert Tangerman. Self-similarity and growth in Birkhoff sums for the golden rotation. *Nonlinearity*, 24:3115–3127, 2011.
- [80] Neal Koblitz. *p-adic numbers, p-adic analysis, and zeta-functions*, volume 58 of *Graduate Texts in Mathematics*. Springer, second edition, 1984.
- [81] J. F. Koksma. *Diophantische Approximationen*, volume 4 of *Ergebnisse der Mathematik und ihrer Grenzgebiete*. Julius Springer, Berlin, 1936.
- [82] Louis Kollross. Prof. Dr. Jérôme Franel, 1859–1939. *Verhandlungen der Schweizerischen Naturforschenden Gesellschaft*, 120:439–444, 1940.
- [83] Ulrich Krengel. *Ergodic theorems*, volume 6 of *De Gruyter Studies in Mathematics*. De Gruyter, 1985. With a Supplement by Antoine Brunei.
- [84] A. H. Kruse. Estimates of $\sum_{k=1}^N k^{-s} \langle kx \rangle^{-t}$. *Acta Arith*, 12:229–261, 1966/1967.
- [85] L. Kuipers and H. Niederreiter. *Uniform distribution of sequences*. John Wiley & Sons, 1974.
- [86] Jeffrey C. Lagarias. Number theory and dynamical systems. In Stefan A. Burr, editor, *The Unreasonable Effectiveness of Number Theory*, volume 46 of *Proceedings of Symposia in Applied Mathematics*, pages 35–72. American Mathematical Society, Providence, RI, 1992.
- [87] Jeffrey C. Lagarias. Number theory zeta functions and dynamical zeta functions. In Thomas Branson, editor, *Spectral Problems in Geometry and Arithmetic*, volume 237 of *Contemporary Mathematics*, pages 45–86. American Mathematical Society, Providence, RI, 1999.
- [88] Matilde Lalín, Francis Rodrigue, and Mathew Rogers. Secant zeta functions. *J. Math. Anal. Appl.*, 409(1):197–204, 2014.
- [89] Edmund Landau. Question 1151. *L’Intermédiaire des mathématiciens*, 8:140–143, 1901. Collected Works, vol. 1, pp. 184–187.
- [90] Edmund Landau. *Handbuch der Lehre von der Verteilung der Primzahlen, erster Band*. B. G. Teubner, Leipzig and Berlin, 1909.
- [91] Edmund Landau. Bemerkungen zu der vorstehenden Abhandlung von Herrn Franel. *Nachrichten von der Gesellschaft der Wissenschaften zu Göttingen, Mathematisch-Physikalische Klasse*, pages 202–206, 1924. Collected Works, vol. 8, pp. 166–170.

- [92] Serge Lang. *Introduction to diophantine approximations*. Addison-Wesley, Reading, MA, 1966.
- [93] Reinhard C. Laubenbacher and David J. Pongelley. Eisenstein’s misunderstood geometric proof of the quadratic reciprocity theorem. *College Math. J.*, 25(1):29–34, 1994.
- [94] Detlef Laugwitz. *Bernhard Riemann 1826–1866: Turning Points in the Conception of Mathematics*. Birkhäuser, 2008. Translated from the German by Abe Shenitzer.
- [95] Franz Lemmermeyer. *Reciprocity laws: from Euler to Eisenstein*. Springer Monographs in Mathematics. Springer, 2000.
- [96] Matyáš Lerch. Question 1547. *L’Intermédiaire des mathématiciens*, 11:144–145, 1904.
- [97] Matyáš Lerch. Sur un série analogue aux fonctions modulaires. *Comptes rendus hebdomadaires des séances de l’Académie des Sciences*, 138:952–954, 1904.
- [98] D. S. Lubinsky. The size of $(q; q)_n$ for q on the unit circle. *J. Number Theory*, 76(2):217–247, 1999.
- [99] S. Marmi and D. Sauzin. *Quasianalytic Monogenic Solutions of a Cohomological Equation*. Number 780 in Memoirs of the American Mathematical Society. American Mathematical Society, Providence, RI, 2003.
- [100] Yves Meyer. *Algebraic Numbers and Harmonic Analysis*, volume 2 of *North-Holland Mathematical Library*. North-Holland Publishing Company, 1972.
- [101] John Milnor. *Dynamics in One Complex Variable*. Number 160 in Annals of Mathematics Studies. Princeton University Press, third edition, 2006.
- [102] Hugh L. Montgomery. *Ten lectures on the interface between analytic number theory and harmonic analysis*. Number 84 in CBMS Regional Conference Series in Mathematics. American Mathematical Society, Providence, RI, 1994.
- [103] Hugh L. Montgomery and Robert C. Vaughan. *Multiplicative Number Theory I: Classical Theory*, volume 97 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, 2006.
- [104] L. J. Mordell. Some aspects of Hardy’s mathematical work: Diophantine approximation. *J. London Math. Soc.*, 25:109–114, 1950.
- [105] A. A. Muromskiii. Na nekotorykh ryadakh po trigonometricheskim funktsiyam. *Izv. Akad. Nauk SSSR Ser. Mat.*, 28(1):53–62, 1964.

- [106] Władysław Narkiewicz. *Rational number theory in the 20th century: from PNT to FLT*. Springer Monographs in Mathematics. Springer, 2012.
- [107] Melvyn B. Nathanson. *Additive number theory: the classical bases*, volume 164 of *Graduate Texts in Mathematics*. Springer, 1996.
- [108] Erwin Neuwenschwander. Riemann’s example of a continuous, ‘nondifferentiable’ function. *Math. Intelligencer*, 1(1):40–44, 1978.
- [109] V. A. Oskolkov. Hardy-Littlewood problems on the uniform distribution of arithmetic progressions. *Math. USSR-Izv.*, 36(1):169–182, 1991.
- [110] Alexander Ostrowski. Bemerkungen zur Theorie der Diophantischen Approximationen. *Abhandlungen aus dem mathematischen Seminar der Hamburgischen Universität*, 1:77–98, 1922. Collected Mathematical Papers, vol. 3, pp. 57–78.
- [111] John C. Oxtoby. *Measure and category: a survey of the analogies between topological and measure spaces*, volume 2 of *Graduate Texts in Mathematics*. Springer, second edition, 1980.
- [112] Alfredo Perna. Elenco delle opere scientifiche del prof. E. Cesàro. *Giornale di matematiche di Battaglini*, 45:320–333, 1907.
- [113] Oskar Perron. *Die Lehre von den Kettenbrüchen*. B. G. Teubner, Leipzig, second edition, 1929.
- [114] S. S. Pillai. On a problem in Diophantine approximation. *Proceedings of the Indian Academy of Sciences, Section A*, 15(3):177–189, 1942.
- [115] S. S. Pillai. On algebraic irrationals. *Proceedings of the Indian Academy of Sciences, Section A*, 15(3):173–176, 1942.
- [116] Mark Pollicott. A note on uniform distribution for primes and closed orbits. *Israel J. Math.*, 55(2):199–212, 1986.
- [117] George Pólya and Gábor Szegő. *Problems and theorems in analysis, volume I*, volume 193 of *Die Grundlehren der mathematischen Wissenschaften*. Springer, 1972. Translated from the German by D. Aepli.
- [118] George Pólya and Gábor Szegő. *Problems and theorems in analysis, volume II*, volume 216 of *Die Grundlehren der mathematischen Wissenschaften*. Springer, 1976. Translated from the German by C. E. Billigheimer.
- [119] Alfred Pringsheim. Grundlagen der allgemeinen Funktionenlehre. In H. Burkhardt, W. Wirtinger, and R. Fricke, editors, *Encyklopädie der mathematischen Wissenschaften mit Einschluss ihrer Anwendungen, Band II, 1. Teil, 1. Hälfte*, pages 1–53. B. G. Teubner, Leipzig, 1899–1916.

- [120] Hervé Queffélec and Martine Queffélec. *Diophantine approximation and Dirichlet series*, volume 2 of *HRI Lecture Notes Series*. Hindustan Book Agency, New Delhi, 2013.
- [121] Quemquaeris. Question 1490. *L'Intermédiaire des mathématiciens*, 6:80, 1899.
- [122] J. L. Raabe. *Die Jacob Bernoullische Function*. Orell, Füssli und Comp., Zürich, 1848.
- [123] Hans Rademacher and Emil Grosswald. *Dedekind sums*. Number 16 in Carus Mathematical Monographs. Mathematical Association of America, 1972.
- [124] Reinhold Remmert. *Theory of Complex Functions*, volume 122 of *Graduate Texts in Mathematics*. Springer, 1991. Translated by Robert B. Burckel.
- [125] Bernhard Riemann. Ueber die Darstellbarkeit einer Function durch eine trigonometrische Reihe. *Abhandlungen der Königlichen Gesellschaft der Wissenschaften zu Göttingen*, 13:87–132, 1866/1867. Gesammelte mathematische Werke, second ed., vol. II, 227–271.
- [126] Tanguy Rivoal. On the convergence of Diophantine Dirichlet series. *Proc. Edinb. Math. Soc. (2)*, 55(2):513–541, 2012.
- [127] Andrew M. Rockett and Peter Szűsz. *Continued fractions*. World Scientific, Singapore, 1992.
- [128] Walter Rudin. *Fourier analysis on groups*. Number 12 in Interscience Tracts in Pure and Applied Mathematics. Interscience Publishers, New York, 1962.
- [129] Walter Rudin. *Principles of mathematical analysis*. International Series in Pure and Applied Mathematics. McGraw-Hill, third edition, 1976.
- [130] Walter Rudin. *Real and Complex Analysis*. McGraw-Hill, third edition, 1987.
- [131] W. H. Schikhof. *Ultrametric calculus: An introduction to p-adic analysis*, volume 4 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, 1984.
- [132] Wolfgang M. Schmidt. Approximation to algebraic numbers. *Enseign. Math. (2)*, 17:187–253, 1971.
- [133] Johannes Schoißengeier. Eine explizite Formel für $\sum_{n \leq N} B_2(\{n\alpha\})$. In Edmund Hlawka, editor, *Zahlentheoretische Analysis, II*, volume 1262 of *Lecture Notes in Mathematics*, pages 134–138. Springer, 1987.

- [134] Sanford L. Segal. *Nine introductions in complex analysis*, volume 208 of *North-Holland Mathematics Studies*. Elsevier, revised edition, 2008.
- [135] Jeffrey Shallit. Real numbers with bounded partial quotients: a survey. *Enseign. Math. (2)*, 38(1-2):151–187, 1992.
- [136] Waclaw Sierpiński. Un théorème sur les nombres irrationnels. *Bulletin international de l'Académie des sciences de Cracovie A*, pages 725–727, 1909. Oeuvres Choiesies, tome I, pp. 155–157.
- [137] Ya. G. Sinai. *Topics in Ergodic Theory*, volume 44 of *Princeton Mathematical Series*. Princeton University Press, 1994.
- [138] Ya. G. Sinai, editor. *Dynamical systems, ergodic theory and applications*, volume 100 of *Encyclopaedia of Mathematical Sciences*. Springer, second edition, 2000.
- [139] Yakov G. Sinai and Corinna Ulcigrai. Estimates from above of certain double trigonometric sums. *J. Fixed Point Theory Appl.*, 6:93–113, 2009.
- [140] Josef Škrášek. Seznam prací prof. Matyáše Lercha. *Časopis pro pěstování matematiky*, 78(2):139–148, 1953.
- [141] David Eugene Smith, editor. *A source book in mathematics, Volume 1*. Dover Publications, 1959.
- [142] D. C. Spencer. On a Hardy-Littlewood problem of diophantine approximation. *Proc. Cambridge Philos. Soc.*, 35(4):527–547, 1939. Selecta, vol. 1, pp. 3–23.
- [143] J. Michael Steele. *The Cauchy-Schwarz master class: an introduction to the art of mathematical inequalities*. Cambridge University Press, 2004.
- [144] E. C. Titchmarsh. *The theory of functions*. Oxford University Press, second edition, 1939.
- [145] Giancarlo Travaglini. *Number Theory, Fourier Analysis and Geometric Discrepancy*, volume 81 of *London Mathematical Society Student Texts*. Cambridge University Press, 2014.
- [146] J. Vaaler. A Tauberian theorem related to Weyl's criterion. In G. Rauzy, editor, *Répartition modulo 1. Actes du Colloque de Marseille-Luminy, 4 au 7 Juin 1974*, volume 475 of *Lecture Notes in Mathematics*, pages 253–258. Springer, 1975.
- [147] R. C. Vaughan. *The Hardy-Littlewood Method*, volume 80 of *Cambridge Tracts in Mathematics*. Cambridge University Press, 1981.
- [148] William A. Veech. Topological dynamics. *Bull. Amer. Math. Soc.*, 83(5):775–830, 1977.

- [149] Paul Verschueren and Ben Mestel. Growth of the Sudler product of sines at the golden rotation number. *J. Math. Anal. Appl.*, 433(1):200–226, 2016.
- [150] I. M. Vinogradov. *The method of trigonometrical sums in the theory of numbers*. Interscience Publishers, 1954. Translated from the Russian by K. F. Roth and Anne Davenport.
- [151] Ilya Vinogradov. A generalization of a result of Hardy and Littlewood, 2011. arXiv:0709.2882.
- [152] Jan von Plato. *Creating Modern Probability: Its Mathematics, Physics and Philosophy in Historical Perspective*. Cambridge Studies in Probability, Induction and Decision Theory. Cambridge University Press, 1994.
- [153] Arnold Walfisz. Über einige trigonometrische Summen. *Math. Z.*, 33:564–601, 1931.
- [154] G. N. Watson. The sum of a series of cosecants. *Philosophical Magazine Series 6*, 31(182):111–118, 1916.
- [155] Derek Thomas Whiteside. Patterns of mathematical thought in the later seventeenth century. *Arch. Hist. Exact Sci.*, 1(3):179–388, 1961.
- [156] W. Ll. G. Williams. The sum of a series of cosecants. *Philosophical Magazine Series 7*, 19(126):402–404, 1935.
- [157] J. R. Wilton. An approximate functional equation with applications to a problem of Diophantine approximation. *J. Reine Angew. Math.*, 169:219–237, 1933.
- [158] Thomas H. Wolff. *Lectures on Harmonic Analysis*, volume 29 of *University Lecture Series*. American Mathematical Society, Providence, RI, 2003.
- [159] Jean-Christophe Yoccoz. Analytic linearization of circle diffeomorphisms. In Hakan Eliasson, Sergei Kuksin, Stefano Marmi, and Jean-Christophe Yoccoz, editors, *Dynamical Systems and Small Divisors*, volume 1784 of *Lecture Notes in Mathematics*, pages 125–173. Springer, 2002.